



Spectral approach for kernel-based interpolation

Bertrand Gauthier, Xavier Bay

► To cite this version:

Bertrand Gauthier, Xavier Bay. Spectral approach for kernel-based interpolation. Annales de la Faculté des Sciences de Toulouse. Mathématiques., 2012, 21 (3), pp.439-479. hal-00915529v3

HAL Id: hal-00915529

<https://hal.science/hal-00915529v3>

Submitted on 13 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SPECTRAL APPROACH FOR KERNEL-BASED INTERPOLATION

Bertrand GAUTHIER & Xavier BAY

Abstract. — We describe how the resolution of a kernel-based interpolation problem can be associated with a spectral problem. An integral operator is defined from the embedding of the considered Hilbert subspace into an auxiliary Hilbert space of square-integrable functions. We finally obtain a spectral representation of the interpolating elements which allows their approximation by spectral truncation. As an illustration, we show how this approach can be used to enforce boundary conditions in kernel-based interpolation models and in what it offers an interesting alternative for dimension reduction.

Résumé (Approche spectrale pour l'interpolation à noyaux). — Nous décrivons comment la résolution d'un problème d'interpolation à noyaux peut être associée à un problème spectral. Un opérateur intégral est défini à partir d'un plongement du sous-espace hilbertien considéré dans un espace de Hilbert auxiliaire composé de fonctions de carré intégrable. On obtient une représentation spectrale des éléments interpolants permettant leur approximation par troncature du spectre. À titre d'exemple, nous montrons comment cette approche peut être utilisée afin d'intégrer des informations de type conditions aux limites dans un modèle d'interpolation et en quoi elle offre une alternative intéressante pour la réduction de dimension.

Contents

1. Introduction.....	2
2. Optimal interpolation in Hilbert subspaces.....	3
3. Problem-adapted integral operators.....	5
4. Representation and approximation of the optimal interpolator...	12
5. Finite Case.....	16
6. Application to Gaussian process models.....	18

Key words and phrases. — kernel-based interpolation, integral operators, Hilbert subspaces, RKHS, Gaussian process models, spectral decomposition, infinite data set, boundary conditions, optimal approximation.

To quote this article. — B. Gauthier and X. Bay, *Spectral approach for kernel-based interpolation*, Annales de la Faculté des Sciences de Toulouse, Tome 21, numéro 3 (2012), p. 439-479.

7. Example of application.....	22
Acknowledgments.....	32
References.....	32

1. Introduction

This work is devoted to the study of kernel-based interpolation methods (see for instance and among others [Wah90, Ste99, BTA04, RW06]). In order to cover a relatively wide class of problems, we consider the general framework of interpolation in a *separated topological real vector space* E . We denote by E' the topological dual of E and by $\langle \cdot, \cdot \rangle_{E, E'}$ the associated duality bracket. For a linear subspace M of E' and $e \in E$, we say that $f \in E$ is an *interpolator of e for M* (or *on M*) if

$$\forall e' \in M, \langle f, e' \rangle_{E, E'} = \langle e, e' \rangle_{E, E'}.$$

In this context, we focus on the two linked kernel-based methods that are *optimal interpolation in Hilbert subspaces of E* (see e.g., [Att92]) and *Gaussian process models* based on the conditioning of zero-mean Gaussian processes with sample paths in E . We consider interpolation problems associated with general sets M , including more particularly the case where M is infinite dimensional (infinite data set). Such a situation may for instance occur when one aims at enforcing boundary values conditions in an interpolating functions problem.

We propose and analyze an overall process which associates the resolution of a kernel-based interpolation problem with the spectral decomposition of an integral operator. This leads to a spectral representation of the solutions of the considered interpolation problem (Theorem 4.1). By applying spectral truncations, we obtain approximations of the interpolating elements which are optimal in a given sense (Proposition 6.4).

From a theoretical point of view, the spectral properties used in this article are well-known and related to extensions of the Mercer's Theorem and Karhunen-Loève's Theorem. From an applied point of view, the use of spectral methods in approximation and learning is not new either. We can for instance quote the article of F. Cucker and S. Smale [CS02], where recalls and discussions concerning Mercer kernels and their applications in learning theory can be found. One can also refer to the works of E. Parzen [Par62] (also mentioned in [BTA04, Section 2.4]), or among others, to the articles [Wal67, Kue71, Raj72]. Also, remark that our approach is based on the embedding of the considered Hilbert subspace into an auxiliary Hilbert space of square-integrable real-valued functions. The various applications and structures we consider can in this sense be compared with the ones appearing in the work of M.Z. Nashed and G. Wahba [NW74]. The main objective of the present article is to give a theoretical description, in the general context of topological vector spaces, of the processes involved in the association of a kernel-based interpolation problem with a spectral problem. We also aim at showing the potential interests of such an approach.

The first part of this article (Section 2) is devoted to the description of optimal interpolation in Hilbert subspaces. In Section 3, we define the notion of *regular embedding adapted to an interpolation problem*. We also show how this embedding defines an integral operator, which is referred to as *problem-adapted*. In Section 4, we use the spectral decomposition of this operator in order to study the initial interpolation problem and its approximation by spectral truncation.

In Section 5, we consider the case where M is finite-dimensional. Explicit calculations are carried out in order to illustrate the use of spectral considerations in this particular case. Section 6 is next devoted to Gaussian process models. The spectral representations considered in the previous sections are extended to the conditioning problem. In particular, we show the *IMSE-optimal* character of the approximation by truncation in this context.

We finally develop (Section 7) a theoretical example of application involving a Hilbert subspace of continuously differentiable real-valued functions on \mathbb{R}^2 . We consider a particular class of kernels and show how to enforce Cauchy-type constraints on a circle (conditions on both values and normal derivatives) in the associated interpolation models. The difference between approximation by truncation and discretization is illustrated.

2. Optimal interpolation in Hilbert subspaces

2.1. Hilbert subspace and RKHS. — The L. Schwartz theory of Hilbert subspaces [Sch64] (or Hilbertian subspaces) is an equivalent formalism for the more widespread theory of reproducing kernel Hilbert spaces (RKHS), introduced by N. Aronszajn in [Aro50], this equivalence is for instance discussed in Remark 2.1. The abstract formalism of L. Schwartz is adapted to the framework of topological vector spaces; it also draws interesting parallels with operator theory (see for instance Proposition 3.7).

2.1.1. Hilbert subspace. — The general framework of the Hilbert subspaces of E requires the (real) topological vector space E to be also locally convex and quasi-complete (see for instance [Rud95, Sch64]), what we assume thereafter. Remark that these properties are verified by most of the classical functions spaces and in particular by all Fréchet and Banach spaces. We denote by E' the topological dual space of E .

A *Hilbert subspace* \mathcal{H} of E is a linear subspace of E endowed with a Hilbert structure such that the inclusion of the Hilbert space \mathcal{H} into E is *continuous*. We use the notation $\mathcal{H} \in \text{Hilb}(E)$. We then denote by $T_{\mathcal{H}}$ the *Hilbert kernel* naturally associated with $\mathcal{H} \in \text{Hilb}(E)$. We remind that $T_{\mathcal{H}} : E' \rightarrow \mathcal{H} \subset E$ is a linear, symmetric and positive application,

$$\text{i.e. } \forall e' \text{ and } f' \in E', \langle T_{\mathcal{H}} e', f' \rangle_{E, E'} = \langle T_{\mathcal{H}} f', e' \rangle_{E, E'} \text{ and } \langle T_{\mathcal{H}} e', e' \rangle_{E, E'} \geq 0.$$

The kernel $T_{\mathcal{H}}$ is in particular characterized by the *representation property*,

$$(2.1) \quad \forall h \in \mathcal{H}, \forall e' \in E', \langle h, e' \rangle_{E, E'} = (h | T_{\mathcal{H}} e')_{\mathcal{H}},$$

where $(\cdot|\cdot)_{\mathcal{H}}$ is the inner product of \mathcal{H} . If $\{h_j : j \in \mathbb{J}\}$ is an orthonormal basis of $\mathcal{H} \in \text{Hilb}(E)$, the Hilbert kernel $T_{\mathcal{H}}$ can be written under the form

$$(2.2) \quad T_{\mathcal{H}} = \sum_{j \in \mathbb{J}} h_j \otimes h_j, \text{ i.e. } \forall e' \in E', T_{\mathcal{H}} e' = \sum_{j \in \mathbb{J}} \langle h_j, e' \rangle_{E, E'} h_j.$$

2.1.2. Reproducing kernel Hilbert space. — A RKHS \mathcal{H} of real-valued functions on a set \mathcal{X} is a Hilbert subspace of $\mathbb{R}^{\mathcal{X}}$ (space of real-valued functions on \mathcal{X}) endowed with the topology of the pointwise convergence (see [Aro50, Sch64, BTA04]).

The *reproducing kernel* $K(\cdot, \cdot)$ of \mathcal{H} is then linked with the Hilbert kernel $T_{\mathcal{H}}$ by the relation, for all x and $y \in \mathcal{X}$,

$$(2.3) \quad K(x, y) = \langle T_{\mathcal{H}} \delta_x, \delta_y \rangle_{E, E'},$$

where δ_x is the evaluation functional for $x \in \mathcal{X}$, that is $\langle f, \delta_x \rangle_{E, E'} = f(x)$ for all $f \in E = \mathbb{R}^{\mathcal{X}}$. This definition for a RKHS (as Hilbert subspace) is therefore exactly equivalent to the more common definition, namely that \mathcal{H} is a Hilbert space of real-valued functions on \mathcal{X} such that, for all $x \in \mathcal{X}$, the linear map $L_x : \mathcal{H} \rightarrow \mathbb{R}, h \mapsto h(x)$, is continuous.

Remark 2.1. — The RKHS theory can first appear to be a particular case of the Hilbert subspaces one. In reality, this two theory are equivalent and only differ by their formalism (see [Sch64, Gau11]). Indeed, one can for instance consider E as a linear subspace of $\mathbb{R}^{E'}$ (since $E \subset E''$) and then assimilate a Hilbert subspace of E with a RKHS of real-valued functions on E' .

2.2. Optimal interpolation. — Let $\mathcal{H} \in \text{Hilb}(E)$ and M be a linear subspace of E' . For a given $\varphi \in \mathcal{H}$, the set of the elements of \mathcal{H} which interpolate φ on M can be easily described thanks to the Hilbert subspace structure of \mathcal{H} . In what follows, we resume some of the main results concerning such a study. Throughout this article, we will frequently refer to the *interpolation problem associated with $\mathcal{H} \in \text{Hilb}(E)$ and M* , without necessarily mention the element of \mathcal{H} which has to be interpolated.

Let us introduce the set

$$M^0 = \left\{ e \in E : \forall e' \in M, \langle e, e' \rangle_{E, E'} = 0 \right\}.$$

We define $\mathcal{H}_0 = M^0 \cap \mathcal{H} = T_{\mathcal{H}}(M)^{\perp}$, where $T_{\mathcal{H}}(M)^{\perp}$ denotes the orthogonal, in \mathcal{H} , of $T_{\mathcal{H}}(M) = \{T_{\mathcal{H}} e' : e' \in M\}$. Then, for a fixed $\varphi \in \mathcal{H}$,

$$\varphi + (M^0 \cap \mathcal{H})$$

is the set of all interpolators, in \mathcal{H} , of φ for M .

$\varphi + (M^0 \cap \mathcal{H})$ is a non-empty closed affine subspace of \mathcal{H} and is therefore also convex. Thus $\varphi + (M^0 \cap \mathcal{H})$ admits a minimal norm element, which we denote $h_{\varphi, M}$ and call *minimal norm interpolator*, or *optimal interpolator*. $h_{\varphi, M}$ is then the orthogonal projection of 0 onto $\varphi + (M^0 \cap \mathcal{H})$. Let us remark that this first characterization of the optimal interpolator is essentially *non-constructive*, in the sense that it does not allow the construction of $h_{\varphi, M}$ from the only knowledge of the values of φ on M .

By definition of the orthogonal projection, $h_{\varphi,M} - 0$ is orthogonal to \mathcal{H}_0 ,

$$\text{i.e. } h_{\varphi,M} \in \mathcal{H}_0^\perp = \left(T_{\mathcal{H}}(M)^\perp \right)^\perp = \overline{T_{\mathcal{H}}(M)}^{\mathcal{H}} = \mathcal{H}_M,$$

with \mathcal{H}_M the closure, in \mathcal{H} , of the linear space spanned by the $T_{\mathcal{H}}e'$, for $e' \in M$. This introduces the orthogonal decomposition $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_M$ and implies in particular that $h_{\varphi,M}$ is the only interpolator, in \mathcal{H}_M , of φ for M .

Finally, let $P_{\mathcal{H}_M}$ be the orthogonal projection of \mathcal{H} onto \mathcal{H}_M . We know that $\varphi - P_{\mathcal{H}_M}[\varphi]$ is orthogonal to \mathcal{H}_M , thus $\varphi - P_{\mathcal{H}_M}[\varphi] \in \mathcal{H}_0$, i.e. $P_{\mathcal{H}_M}[\varphi]$ interpolates φ for M . We finally obtain

$$h_{\varphi,M} = P_{\mathcal{H}_M}[\varphi]$$

and this second characterization is suitable for the construction of $h_{\varphi,M}$ from the only knowledge of $\langle \varphi, e' \rangle_{E,E'}$, for $e' \in M$.




FIGURE 1. Schematic representation of optimal interpolation in a Hilbert subspace.

The Hilbert kernel $T_{\mathcal{H}_M}$ of the Hilbert subspace \mathcal{H}_M , $(\cdot|\cdot)_{\mathcal{H}}$, is linked with $T_{\mathcal{H}}$ by the relation

$$T_{\mathcal{H}_M} = P_{\mathcal{H}_M} T_{\mathcal{H}}.$$

Hence, the knowledge of $T_{\mathcal{H}_M}$ defines the orthogonal projection $P_{\mathcal{H}_M}$ and reciprocally, this result staying true for any closed linear subspace of \mathcal{H} . This implies in particular that the Hilbert kernel $T_{\mathcal{H}_0}$ of \mathcal{H}_0 , $(\cdot|\cdot)_{\mathcal{H}}$, is given by $T_{\mathcal{H}_0} = T_{\mathcal{H}} - T_{\mathcal{H}_M}$.

3. Problem-adapted integral operators

Let $\mathcal{H} \in \text{Hilb}(E)$ and let M be a linear subspace of E' . Throughout this section, we consider the interpolation problem associated with \mathcal{H} and M . We use the same notations and definitions as Section 2. Let us in particular remind the orthogonal decomposition $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_M$.

In what follows, we introduce the notion of *regular embeddings associated with an interpolation problem* and study the integral operators they naturally define. This

leads to the construction of specific orthonormal bases of \mathcal{H}_M which are (in the sense of equation (3.14)) suitable for the resolution of the considered interpolation problem.

The results of Sections 3 and 4 hold for any Hilbert subspace \mathcal{H} of E , separable or non-separable. However, if \mathcal{H} is non-separable, the existence of a regular embedding requires \mathcal{H}_M to be separable (see Remark 3.4). Let us mention that complementary considerations concerning Section 3 can be found in [Gau11].

3.1. Regular embedding and parameterization. — Let $(\mathcal{S}, \mathcal{A}, \nu)$ be a general measured set with ν a σ -finite measure. We denote by $L^2(\mathcal{S}, \nu)$ the Hilbert space of square-integrable real-valued functions on \mathcal{S} with respect to ν . Let us remind that $L^2(\mathcal{S}, \nu)$ is in fact a quotient space. Nevertheless, we make the widespread abuse of notation which consists in assimilating elements of $L^2(\mathcal{S}, \nu)$ with functions on \mathcal{S} (instead of considering *equivalence classes* of ν -almost everywhere equal functions). We will sometime refer to $L^2(\mathcal{S}, \nu)$ as the *auxiliary Hilbert space*.

Let $(\cdot|\cdot)_{L^2}$ and $\|\cdot\|_{L^2}$ be respectively the inner product and the norm of $L^2(\mathcal{S}, \nu)$. We recall that

$$\forall f \text{ and } g \in L^2(\mathcal{S}, \nu), (f|g)_{L^2} = \int_{\mathcal{S}} f(s)g(s)d\nu(s).$$

Let us consider an application $\gamma : \mathcal{S} \rightarrow E'$. For all $h \in \mathcal{H}$, we then define the function

$$(3.1) \quad \mathfrak{F}h : \mathcal{S} \rightarrow \mathbb{R} \text{ with } \mathfrak{F}h(s) = \langle h, \gamma s \rangle_{E, E'} \text{ for all } s \in \mathcal{S}.$$

We now introduce conditions concerning $L^2(\mathcal{S}, \nu)$, γ and \mathcal{H} , namely:

- C-i. for all $h \in \mathcal{H}$, the function $\mathfrak{F}h : \mathcal{S} \rightarrow \mathbb{R}$ is measurable,
- C-ii. the function $(s, t) \in \mathcal{S} \times \mathcal{S} \mapsto \langle T_{\mathcal{H}}\gamma s, \gamma t \rangle_{E, E'} = (T_{\mathcal{H}}\gamma s | T_{\mathcal{H}}\gamma t)_{\mathcal{H}}$ is measurable,
- C-iii. $\tau = \int_{\mathcal{S}} \|T_{\mathcal{H}}\gamma s\|_{\mathcal{H}}^2 d\nu(s) < +\infty$.

Proposition 3.1. — *Under Conditions C-i, C-ii and C-iii, we have $\mathfrak{F}h \in L^2(\mathcal{S}, \nu)$ for all $h \in \mathcal{H}$ and*

$$(3.2) \quad \|\mathfrak{F}h\|_{L^2}^2 \leq \tau \|h\|_{\mathcal{H}}^2.$$

Hence, the linear application $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$, $h \mapsto \mathfrak{F}h$ is continuous.

Proof. — Representation property (2.1), Cauchy-Schwarz inequality applied to $(\cdot|\cdot)_{\mathcal{H}}$ and finally Condition C-iii give

$$(3.3) \quad \int_{\mathcal{S}} \langle h, \gamma s \rangle_{E, E'}^2 d\nu(s) = \int_{\mathcal{S}} (h | T_{\mathcal{H}}\gamma s)_{\mathcal{H}}^2 d\nu(s) \leq \tau \|h\|_{\mathcal{H}}^2,$$

each integral being well-defined thanks to Conditions C-i and C-ii. \square

We now consider the orthogonal decomposition $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_M$ and add the following condition on the application $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$,

- C-iv. for all $h \in \mathcal{H}$, $\|\mathfrak{F}h\|_{L^2} = 0$ if and only if $h \in \mathcal{H}_0$.

Definition 3.2. — We call *regular embedding of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$ adapted to the interpolation problem associated with \mathcal{H} and M* an application $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$ defined from a *parameterization* $\gamma : \mathcal{S} \rightarrow E'$ via equation (3.1) and which verifies Conditions C-i, C-ii, C-iii and C-iv.

Let $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$ be a regular embedding of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$. We consider the linear subspace $\mathfrak{F}(\mathcal{H})$ of $L^2(\mathcal{S}, \nu)$ given by $\mathfrak{F}(\mathcal{H}) = \{\mathfrak{F}h : h \in \mathcal{H}\}$ (the image of \mathcal{H} through \mathfrak{F}). From C-iv, $\mathfrak{F}(\mathcal{H}_0) = 0$, hence $\mathfrak{F}(\mathcal{H}) = \mathfrak{F}(\mathcal{H}_M)$. We endow this space of the following inner-product,

$$(3.4) \quad \forall h \text{ and } g \in \mathcal{H}_M, (\mathfrak{F}h | \mathfrak{F}g)_{\mathfrak{F}(\mathcal{H})} = (h | g)_{\mathcal{H}}.$$

Proposition 3.3. — Let $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$ be a regular embedding of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$, then $\mathfrak{F}(\mathcal{H}_M)$, $(\cdot | \cdot)_{\mathfrak{F}(\mathcal{H})}$ is a Hilbert space. It is isometric to \mathcal{H}_M , $(\cdot | \cdot)_{\mathcal{H}}$, the isometry being the restriction to \mathcal{H}_M of the regular embedding \mathfrak{F} .

In addition, the inclusion of the Hilbert space $\mathfrak{F}(\mathcal{H}_M)$, $(\cdot | \cdot)_{\mathfrak{F}(\mathcal{H})}$ into $L^2(\mathcal{S}, \nu)$ is continuous. In other words, $\mathfrak{F}(\mathcal{H}_M) \in \text{Hilb}(L^2(\mathcal{S}, \nu))$.

Proof. — The fact that $\mathfrak{F}(\mathcal{H}_M)$ is a Hilbert space isometric to \mathcal{H}_M directly follows from its construction. Further, from Proposition 3.1, we have for all $h \in \mathcal{H}_M$,

$$\|\mathfrak{F}h\|_{L^2}^2 \leq \tau \|h\|_{\mathcal{H}}^2 = \tau \|\mathfrak{F}h\|_{\mathfrak{F}(\mathcal{H})}^2,$$

thus $\mathfrak{F}(\mathcal{H}_M) \in \text{Hilb}(L^2(\mathcal{S}, \nu))$. □

Remark 3.4. — One can for instance consult [For85] for a discussion about the conditions appearing in Definition 3.2. The ones we use here are of similar type but specially adapted to the study of the interpolation problem associated with \mathcal{H} and M . Indeed, let \mathfrak{F} be a regular embedding of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$ and consider

$$(3.5) \quad \int_{\mathcal{S}} f(s) \langle h, \gamma s \rangle_{E, E'} d\nu(s) = (f | \mathfrak{F}h)_{L^2},$$

where f is a fixed element of $L^2(\mathcal{S}, \nu)$ and $h \in \mathcal{H}$. From Condition C-iv, the value of expression (3.5) does not vary if we replace h by $h + h_0$, with $h_0 \in \mathcal{H}_0$. This quantity thus only depends of the values of h on M (i.e. of $\langle h, e' \rangle_{E, E'}$ for $e' \in M$), which are the only available informations when considering an interpolation problem associated with M and an element h of \mathcal{H} .

When \mathcal{S} is a topological space (endowed with its Borel σ -algebra) and Conditions C-i, C-ii and C-iii are already verified, C-iv will for instance be realized if for all $h \in \mathcal{H}$, the functions $\mathfrak{F}h$ are continuous and if $M = \text{span}\{\gamma(\text{supp}(\nu))\}$ (i.e. M is the linear subspace of E' spanned by $\gamma(\text{supp}(\nu))$, with $\text{supp}(\nu)$ the support of ν).

Finally, note that the existence of a regular embedding \mathfrak{F} associated with the interpolation problem defined by \mathcal{H} and M implies in particular that \mathcal{H}_M is separable, see for instance Proposition 3.8.

Example 3.5. — Let us consider a RKHS \mathcal{H} of continuous real-valued functions on a topological space \mathcal{X} and the problem consisting in the interpolation of an element φ of \mathcal{H} at given points x_1, \dots, x_n of \mathcal{X} (i.e. $M = \text{span}\{\delta_{x_1}, \dots, \delta_{x_n}\}$).

One can for instance define a regular embedding for this problem by introducing the measure $\nu = \sum_{i=1}^n w_i \delta_{x_i}$ (with $w_i > 0$) on $\mathcal{S} = \mathcal{X}$ (endowed with its Borel σ -algebra) and the parameterization $\gamma : x \mapsto \delta_x$ (another possible parameterization for this problem is given in Section 5).

More generally, if we suppose that the values of φ are known on a closed subset D of \mathcal{X} (*i.e.* $M = \text{span}\{\delta_x, x \in D\}$) while keeping the same parameterization γ , one just has to consider a measure ν on \mathcal{X} whose support is D and such that Conditions C-iii is also verified.

Note that in this particular example, if one identifies elements of $L^2(\mathcal{S}, \nu)$ with functions on $\mathcal{S} = \mathcal{X}$, then the application \mathfrak{F} is in fact the identity operator on \mathcal{H} , with $\mathfrak{F}h(x) = h(x)$, for all $h \in \mathcal{H}$ and $x \in \mathcal{X}$.

Proposition 3.6. — *Let $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$ be a regular embedding of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$ and consider its adjoint operator ${}^t\mathfrak{F} : L^2(\mathcal{S}, \nu) \rightarrow \mathcal{H}$ defined by equation (3.7) hereafter. Then, for all $f \in L^2(\mathcal{S}, \nu)$, ${}^t\mathfrak{F}f \in \mathcal{H}_M$ and we have the following integral representation,*

$$(3.6) \quad \forall f \in L^2(\mathcal{S}, \nu), {}^t\mathfrak{F}f = \int_{\mathcal{S}} f(s) T_{\mathcal{H}} \gamma s \, d\nu(s),$$

this expression having to be understood in the sense of equation (3.8).

Proof. — We remind that ${}^t\mathfrak{F} : L^2(\mathcal{S}, \nu) \rightarrow \mathcal{H}$ is defined by

$$(3.7) \quad \forall h \in \mathcal{H}, \forall f \in L^2(\mathcal{S}, \nu), (h | {}^t\mathfrak{F}f)_{\mathcal{H}} = (\mathfrak{F}h | f)_{L^2}.$$

From C-iv, we directly deduce that ${}^t\mathfrak{F}f \in \mathcal{H}_M$ for all $f \in L^2(\mathcal{S}, \nu)$. Next, by applying the preceding equation to $h = T_{\mathcal{H}} e'$ with $e' \in E'$, we obtain

$$(3.8) \quad \begin{aligned} \langle {}^t\mathfrak{F}f, e' \rangle_{E, E'} &= \int_{\mathcal{S}} f(s) \langle T_{\mathcal{H}} e', \gamma s \rangle_{E, E'} \, d\nu(s) \\ &= \int_{\mathcal{S}} f(s) \langle T_{\mathcal{H}} \gamma s, e' \rangle_{E, E'} \, d\nu(s), \end{aligned}$$

which corresponds to equation (3.6) (one can refer to [Bou59] for details concerning the notion of vectorial integral). \square

3.2. Integral operator defined by a regular embedding. — We still consider the same interpolation problem associated with \mathcal{H} and M . Thanks to Proposition 3.3, we know that a regular embedding \mathfrak{F} of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$ defines a Hilbert subspace $\mathfrak{F}(\mathcal{H}_M)$, $(\cdot | \cdot)_{\mathfrak{F}(\mathcal{H}_M)}$ of $L^2(\mathcal{S}, \nu)$. Hence, from the Hilbert subspaces theory, it admits a unique associated Hilbert kernel. If we identify the continuous dual of $L^2(\mathcal{S}, \nu)$ with itself (Riesz-Fréchet representation Theorem), the Hilbert kernel of $\mathfrak{F}(\mathcal{H}_M)$ relatively to $L^2(\mathcal{S}, \nu)$ is the unique linear application

$$\mathcal{L}_{\gamma, \nu} : (L^2(\mathcal{S}, \nu))' = L^2(\mathcal{S}, \nu) \rightarrow \mathfrak{F}(\mathcal{H}_M) \subset L^2(\mathcal{S}, \nu)$$

which verifies the representation property, for all $h \in \mathcal{H}$ and $f \in L^2(\mathcal{S}, \nu)$,

$$(3.9) \quad (\mathfrak{F}h | f)_{L^2} = (\mathfrak{F}h | \mathcal{L}_{\gamma, \nu}[f])_{\mathfrak{F}(\mathcal{H}_M)}.$$

Proposition 3.7. — Let $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$ be a regular embedding of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$ and let $\mathcal{L}_{\gamma, \nu}$ be the kernel of the Hilbert subspace $\mathfrak{F}(\mathcal{H}_M)$ of $L^2(\mathcal{S}, \nu)$, then $\mathcal{L}_{\gamma, \nu} = \mathfrak{F}^t \mathfrak{F}$, i.e. for all $t \in \mathcal{S}$ and $f \in L^2(\mathcal{S}, \nu)$,

$$(3.10) \quad \mathcal{L}_{\gamma, \nu}[f](t) = \int_{\mathcal{S}} (T_{\mathcal{H}} \gamma s | T_{\mathcal{H}} \gamma t)_{\mathcal{H}} f(s) d\nu(s).$$

Proof. — By combining equations (3.9), (3.7) and (3.4), we obtain that, for $h \in \mathcal{H}$ and $f \in L^2(\mathcal{S}, \nu)$,

$$(\mathfrak{F}h|f)_{L^2} = (h|{}^t\mathfrak{F}f)_{\mathcal{H}} = (\mathfrak{F}h|\mathfrak{F}^t\mathfrak{F}f)_{\mathfrak{F}(\mathcal{H})} = (\mathfrak{F}h|\mathcal{L}_{\gamma, \nu}[f])_{\mathfrak{F}(\mathcal{H})}.$$

We finally deduce equation (3.10) from the integral expression of ${}^t\mathfrak{F}$ given in Proposition 3.6 (more precisely, equation (3.6)) by applying the preceding relation to $h = T_{\mathcal{H}} \gamma t \in \mathcal{H}$, with $t \in \mathcal{S}$. \square

Let us remark that the Hilbert subspace $\mathfrak{F}(\mathcal{H}_M)$ of $L^2(\mathcal{S}, \nu)$ can be assimilated to the RKHS of real-valued functions on \mathcal{S} associated with the reproducing kernel

$$(3.11) \quad \forall (s, t) \in \mathcal{S} \times \mathcal{S}, \mathcal{K}(s, t) = (T_{\mathcal{H}} \gamma t | T_{\mathcal{H}} \gamma s)_{\mathcal{H}}.$$

Hence, $\mathcal{L}_{\gamma, \nu}$ can be seen as a classic integral operator on $L^2(\mathcal{S}, \nu)$ defined by the symmetric and positive kernel $\mathcal{K}(\cdot, \cdot)$ on $\mathcal{S} \times \mathcal{S}$ (see for instance [Sch79, §10] and [For85])

We deduce from the theory of integral operators that $\mathcal{L}_{\gamma, \nu}$ is a *Hilbert-Schmidt operator* and therefore a *compact operator*. So $\mathcal{L}_{\gamma, \nu} : L^2(\mathcal{S}, \nu) \rightarrow L^2(\mathcal{S}, \nu)$ is diagonalizable and its eigenvalues are positive. We denote by λ_i those eigenvalues (repeated according to their algebraic multiplicity) and by $\tilde{\phi}_i \in L^2(\mathcal{S}, \nu)$ the associated eigenfunctions, with $i \in \mathbb{I}$ (and where \mathbb{I} is a general index set). We remind that $\{\tilde{\phi}_i : i \in \mathbb{I}\}$ forms an orthonormal basis of $L^2(\mathcal{S}, \nu)$ and that the set of all *strictly positive* eigenvalues is at most countable.

Proposition 3.8. — Let $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$ be a regular embedding of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$ and consider its associated integral operator

$$\mathcal{L}_{\gamma, \nu} = \mathfrak{F}^t \mathfrak{F} : L^2(\mathcal{S}, \nu) \rightarrow \mathfrak{F}(\mathcal{H}_M) \subset L^2(\mathcal{S}, \nu).$$

Denote by $\{\lambda_n : n \in \mathbb{I}_+\}$ the at most countable set (i.e. $\mathbb{I}_+ \subset \mathbb{N}$) of its strictly positive eigenvalues (repeated according to their multiplicity) and let $\tilde{\phi}_n \in L^2(\mathcal{S}, \nu)$ be their associated eigenfunctions. For all $n \in \mathbb{I}_+$, we define

$$(3.12) \quad \phi_n = \frac{1}{\lambda_n} {}^t\mathfrak{F} \tilde{\phi}_n = \frac{1}{\lambda_n} \int_{\mathcal{S}} \tilde{\phi}_n(s) T_{\mathcal{H}} \gamma s d\nu(s) \in \mathcal{H}_M.$$

Then $\{\sqrt{\lambda_n} \phi_n : n \in \mathbb{I}_+\}$ is an orthonormal basis of the Hilbert space \mathcal{H}_M endowed with the inner product $(\cdot | \cdot)_{\mathcal{H}}$.

Proof. — First remark that from Proposition 3.6, the elements ϕ_n of \mathcal{H}_M are well-defined. By definition, we have that, for all $n \in \mathbb{I}_+$ and for all $h \in \mathcal{H}$,

$$(3.13) \quad (\phi_n | h)_{\mathcal{H}} = \frac{1}{\lambda_n} ({}^t\mathfrak{F} \tilde{\phi}_n | h)_{\mathcal{H}} = \frac{1}{\lambda_n} (\tilde{\phi}_n | \mathfrak{F}h)_{L^2}.$$

As for all $m \in \mathbb{I}_+$, $\mathfrak{F}\phi_m = \tilde{\phi}_m$, the preceding equation (3.13) applied to $h = \phi_m$, gives that $\{\sqrt{\lambda_n}\phi_n : n \in \mathbb{I}_+\}$ is an orthonormal system of \mathcal{H}_M .

From Proposition 3.7 and the properties of Hilbert kernels, we know that the linear subspace spanned by the $\mathcal{L}_{\gamma,\nu}[f]$, $f \in L^2(\mathcal{S}, \nu)$, is dense in $\mathfrak{F}(\mathcal{H}_M)$, $(\cdot|\cdot)_{\mathfrak{F}(\mathcal{H})}$ (and in particular $\{\sqrt{\lambda_n}\tilde{\phi}_n : n \in \mathbb{I}_+\}$ is one of its orthonormal bases). Hence, by isometry between $\mathfrak{F}(\mathcal{H}_M)$, $(\cdot|\cdot)_{\mathfrak{F}(\mathcal{H})}$ and \mathcal{H}_M , $(\cdot|\cdot)_{\mathcal{H}}$, the linear space $\text{span}\{\sqrt{\lambda_n}\phi_n : n \in \mathbb{I}_+\}$ is dense in $\mathfrak{F}(\mathcal{H})$, which concludes the proof. \square

In our context of interpolation, the main interest of the elements ϕ_n of \mathcal{H} , $n \in \mathbb{I}_+$, appearing in Proposition 3.8 is that (see equation (3.13)),

$$(3.14) \quad \forall h \in \mathcal{H}, (\phi_n|h)_{\mathcal{H}} = \frac{1}{\lambda_n} \int_{\mathcal{S}} \tilde{\phi}_n(s) \langle h, \gamma s \rangle_{E,E'} d\nu(s).$$

So, as for equation (3.5) of Remark 3.4, the evaluation of the inner product $(\phi_n|h)_{\mathcal{H}}$ can be directly obtained from the only knowledge of the values of h on M .

We are now able to formulate our representation Theorem 4.1, which simply consists in the use of this particular orthonormal basis and of equation (3.14) in order to describe the orthogonal projection of \mathcal{H} onto \mathcal{H}_M .

Before this, we conclude this section by some additional remarks concerning the structures and applications we have introduced. This will be useful for the rest of our study.

3.3. Some important remarks. — The definition of a regular embedding \mathfrak{F} of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$ allows the construction of various applications and structures in addition to the ones studied until now. This section aims at introducing a few of them. Let us mention the article [NW74], where a similar framework is studied.

3.3.1. Operator on \mathcal{H} defined by a regular embedding. — In the same way as an embedding $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$ defines an integral operator $\mathcal{L}_{\gamma,\nu} = \mathfrak{F}^t \mathfrak{F}$ on $L^2(\mathcal{S}, \nu)$ (see Proposition 3.7), it also defines an operator on \mathcal{H} .

Proposition 3.9. — *Let $\mathfrak{F} : \mathcal{H} \rightarrow L^2(\mathcal{S}, \nu)$ be a regular embedding of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$ and consider the framework of Proposition 3.8. We define the following linear operator on \mathcal{H} ,*

$$(3.15) \quad \forall h \in \mathcal{H}, L_{\gamma,\nu}[h] = {}^t \mathfrak{F} \mathfrak{F} h = \int_{\mathcal{S}} \langle h, \gamma s \rangle_{E,E'} T_{\mathcal{H}} \gamma s d\nu(s).$$

$L_{\gamma,\nu}$ is a continuous symmetric and positive Hilbert-Schmidt operator on the Hilbert space \mathcal{H} , $(\cdot|\cdot)_{\mathcal{H}}$. It is diagonalizable, the eigenspace associated with the null eigenvalues is \mathcal{H}_0 and $\sqrt{\lambda_n}\phi_n$, $n \in \mathbb{I}_+$, are the eigenvectors (with $\|\sqrt{\lambda_n}\phi_n\|_{\mathcal{H}} = 1$) associated with the eigenvalues λ_n , $n \in \mathbb{I}_+$.

Proof. — The properties of symmetry and positivity of $L_{\gamma,\nu}$ are obvious. Let us give a direct proof of the fact that it is a Hilbert-Schmidt operator. Let $\{h_j : j \in \mathbb{J}\}$ be an

orthonormal basis of \mathcal{H} . Using equation (2.2) and Fubini's Theorem, we obtain,

$$(3.16) \quad \sum_{j \in \mathbb{J}} \|L_{\gamma, \nu}[h_j]\|_{\mathcal{H}}^2 = \int_{\mathcal{S}} \int_{\mathcal{S}} (T_{\mathcal{H}} \gamma s | T_{\mathcal{H}} \gamma t)_{\mathcal{H}}^2 d\nu(s) d\nu(t) \leq \tau^2,$$

the last inequality being a consequence of the Cauchy-Schwarz inequality applied to the inner product of \mathcal{H} and of Condition C-iii.

For all $h_0 \in \mathcal{H}_0$, we obviously have ${}^t \mathfrak{F} \mathfrak{F} h_0 = 0$. In addition, for $h \in \mathcal{H}$ and $n \in \mathbb{I}_+$,

$$(3.17) \quad ({}^t \mathfrak{F} \mathfrak{F} \phi_n | h)_{\mathcal{H}} = \lambda_n \left(\frac{1}{\lambda_n} {}^t \mathfrak{F} \tilde{\phi}_n | h \right)_{\mathcal{H}} = (\lambda_n \phi_n | h)_{\mathcal{H}}.$$

This equation combined with Proposition 3.8 completes the spectral decomposition of $L_{\gamma, \nu}$ and also proves its continuity. \square

3.3.2. Two additional Hilbert structures. — We introduce two Hilbert spaces $\overline{\mathfrak{F}(\mathcal{H})}^{L^2}$ and $\overline{\mathcal{H}_M}^{\gamma, \nu}$ which naturally appear when considering a regular embedding \mathfrak{F} of \mathcal{H}_M . Note that these two structures will be useful to us for the application of our approach to Gaussian processes conditioning in Section 6.2.

$\overline{\mathfrak{F}(\mathcal{H})}^{L^2}$ is the closure in $L^2(\mathcal{S}, \nu)$ of the linear subspace $\mathfrak{F}(\mathcal{H})$. Let us notice that $\{\tilde{\phi}_n, n \in \mathbb{I}_+\}$ is obviously one of its orthonormal bases for the inner product $(\cdot | \cdot)_{L^2}$.

Let us now define $\overline{\mathcal{H}_M}^{\gamma, \nu}$. We start by introducing the following symmetric and positive bilinear form on \mathcal{H} , for all h and $g \in \mathcal{H}$,

$$(3.18) \quad (h | g)_{\gamma, \nu} = (\mathfrak{F} h | \mathfrak{F} g)_{L^2} = \int_{\mathcal{S}} \langle h, \gamma s \rangle_{E, E'} \langle g, \gamma s \rangle_{E, E'} d\nu(s).$$

We also set $\|h\|_{\gamma, \nu}^2 = (h | h)_{\gamma, \nu}$. Condition C-iv implies that the nullspace of $(\cdot | \cdot)_{\gamma, \nu}$ is \mathcal{H}_0 (i.e. for $h \in \mathcal{H}$, $\|h\|_{\gamma, \nu} = 0$ if and only if $h \in \mathcal{H}_0$) and \mathcal{H}_M endowed with $(\cdot | \cdot)_{\gamma, \nu}$ is hence a pre-Hilbert space. We then denote by $\overline{\mathcal{H}_M}^{\gamma, \nu}$ the completed of \mathcal{H}_M for $\|\cdot\|_{\gamma, \nu}$.

Remark that the operator $L_{\gamma, \nu}$ (considered as an operator on \mathcal{H}_M) can be naturally extended to $\overline{\mathcal{H}_M}^{\gamma, \nu}$ by continuity. $L_{\gamma, \nu}$ is then a Hilbert-Schmidt operator on $\overline{\mathcal{H}_M}^{\gamma, \nu}$, $(\cdot | \cdot)_{\gamma, \nu}$. It is symmetric and positive definite, its eigenvalues are λ_n , $n \in \mathbb{I}_+$ and each one is associated with the eigenvector ϕ_n (and $\|\phi_n\|_{\gamma, \nu} = 1$).

3.3.3. Isometries. — We are finally in presence of four isometric Hilbert spaces,

$$\mathcal{H}_M, \mathfrak{F}(\mathcal{H}), \overline{\mathfrak{F}(\mathcal{H})}^{L^2} \text{ and } \overline{\mathcal{H}_M}^{\gamma, \nu}.$$

As we have seen in Proposition 3.3, the isometry between \mathcal{H}_M , $(\cdot | \cdot)_{\mathcal{H}}$ and $\mathfrak{F}(\mathcal{H})$, $(\cdot | \cdot)_{\mathfrak{F}(\mathcal{H})}$ is the restriction of \mathfrak{F} to \mathcal{H}_M . The continuous extension of this first isometry defines the isometry between $\overline{\mathcal{H}_M}^{\gamma, \nu}$, $(\cdot | \cdot)_{\gamma, \nu}$ and $\overline{\mathfrak{F}(\mathcal{H})}^{L^2}$, $(\cdot | \cdot)_{L^2}$.

The isometry between $\overline{\mathfrak{F}(\mathcal{H})}^{L^2}$, $(\cdot | \cdot)_{L^2}$ and $\mathfrak{F}(\mathcal{H})$, $(\cdot | \cdot)_{\mathfrak{F}(\mathcal{H})}$ is given by

$$\forall n \in \mathbb{I}_+, \tilde{\phi}_n \rightarrow \sqrt{\lambda_n} \tilde{\phi}_n.$$

It is in fact the restriction of the square-root of $\mathcal{L}_{\gamma,\nu}$ to $\overline{\mathfrak{F}(\mathcal{H})}^{L^2}$, with

$$\mathcal{L}_{\gamma,\nu}^{\frac{1}{2}} \left[\sum_{i \in \mathbb{I}} \alpha_i \tilde{\phi}_i \right] = \sum_{i \in \mathbb{I}} \alpha_i \sqrt{\lambda_i} \tilde{\phi}_i,$$

where $\sum_{i \in \mathbb{I}} \alpha_i \tilde{\phi}_i \in L^2(\mathcal{S}, \nu)$. We obviously have $\mathcal{L}_{\gamma,\nu} = \mathcal{L}_{\gamma,\nu}^{\frac{1}{2}} \circ \mathcal{L}_{\gamma,\nu}^{\frac{1}{2}}$.

3.3.4. Pseudoinverse of a regular embedding. — Let us consider the framework of Proposition 3.8. We can define the pseudoinverse (or generalized inverse) \mathfrak{F}^\dagger of \mathfrak{F} by

$$(3.19) \quad \forall n \in \mathbb{I}_+, \mathfrak{F}^\dagger \tilde{\phi}_n = \phi_n = \frac{1}{\lambda_n} {}^t \mathfrak{F} \tilde{\phi}_n$$

and, for $i \in \mathbb{I} \setminus \mathbb{I}_+$ (i.e. $\lambda_i = 0$), $\mathfrak{F}^\dagger \tilde{\phi}_i = 0$. Then, \mathfrak{F}^\dagger is well-defined from $L^2(\mathcal{S}, \nu)$ onto $\overline{\mathcal{H}_M}^{\gamma,\nu}$ and

$$(3.20) \quad \forall f \in L^2(\mathcal{S}, \nu), \mathfrak{F}^\dagger f = \sum_{n \in \mathbb{I}_+} \left(f \Big| \tilde{\phi}_n \right)_{L^2} \phi_n \in \overline{\mathcal{H}_M}^{\gamma,\nu}.$$

The restriction of \mathfrak{F}^\dagger to $\overline{\mathfrak{F}(\mathcal{H})}^{L^2}$ defines the inverse of the isometry between $\overline{\mathcal{H}_M}^{\gamma,\nu}$ and $\overline{\mathfrak{F}(\mathcal{H})}^{L^2}$. In the same way, its restriction to \mathcal{H}_M gives the inverse of the isometry between \mathcal{H}_M and $\mathfrak{F}(\mathcal{H})$. We have in particular

$$(3.21) \quad P_{\mathcal{H}_M} = \mathfrak{F}^\dagger \mathfrak{F},$$

which is in fact an equivalent formulation of Theorem 4.1.

4. Representation and approximation of the optimal interpolator

For $\mathcal{H} \in \text{Hilb}(E)$, we consider the optimal interpolation problem in \mathcal{H} defined by $\varphi \in \mathcal{H}$ and a linear subspace M of E' . In order to apply Section 3 results, we suppose M such that \mathcal{H}_M is separable (thus, M can be an arbitrary linear subspace of E' if \mathcal{H} is itself separable).

4.1. Spectral representation for optimal interpolation. — Once an orthonormal basis of \mathcal{H}_M known, one can easily express the orthogonal projection of \mathcal{H} onto \mathcal{H}_M . Then, in order to compute the optimal interpolator of $\varphi \in \mathcal{H}$ for M (see Section 2), we need to be able to evaluate the inner-product in \mathcal{H} between φ and each elements of the considered basis of \mathcal{H}_M , and this from the only knowledge of the values of φ on M (which are, in our context of interpolation, the only available informations concerning φ). This is precisely the property of the orthonormal basis of \mathcal{H}_M associated with a regular embedding \mathfrak{F} , its elements indeed verify equation (3.14).

Remark that in order to be applied to a given interpolation problem (associated with \mathcal{H} and M), our approach requires the preliminary choice of a measurable space $(\mathcal{S}, \mathcal{A}, \nu)$ and of a parameterization $\gamma : \mathcal{S} \rightarrow E'$ allowing the definition of a regular embedding \mathfrak{F} of \mathcal{H}_M into the auxiliary space $L^2(\mathcal{S}, \nu)$. Some considerations concerning this choice are discussed in Section 4.3.

Theorem 4.1. — Let \mathfrak{F} be a regular embedding of \mathcal{H}_M into $L^2(\mathcal{S}, \nu)$ and consider the orthonormal basis $\{\sqrt{\lambda_n}\phi_n : n \in \mathbb{I}_+\}$ of \mathcal{H}_M associated with \mathfrak{F} . Then, for $\varphi \in \mathcal{H}$, we have

$$(4.1) \quad P_{\mathcal{H}_M}[\varphi] = \sum_{n \in \mathbb{I}_+} \phi_n \int_{\mathcal{S}} \langle \phi_n, \gamma s \rangle_{E, E'} \langle \varphi, \gamma s \rangle_{E, E'} d\nu(s).$$

Proof. — We just have to consider Proposition 3.8 and equation (3.13),

$$\begin{aligned} P_{\mathcal{H}_M}[\varphi] &= \sum_{n \in \mathbb{I}_+} \sqrt{\lambda_n} \phi_n \left(\sqrt{\lambda_n} \phi_n \Big| \varphi \right)_{\mathcal{H}} = \sum_{n \in \mathbb{I}_+} \lambda_n \phi_n \left(\tilde{\phi}_n \Big| \mathfrak{F} \varphi \right)_{\mathfrak{F}(\mathcal{H})} \\ &= \sum_{n \in \mathbb{I}_+} \phi_n \left(\tilde{\phi}_n \Big| \mathfrak{F} \varphi \right)_{L^2}. \end{aligned}$$

See also equation (3.21) for an equivalent formulation of this result. \square

Let us remark that the sum appearing in equation (4.1) converges by construction in \mathcal{H} . Since $\mathcal{H} \in \text{Hilb}(E)$, it also converges for the initial topology of E and for its weak topology $\sigma(E, E')$. Then, in particular, for all $e' \in E'$,

$$\langle P_{\mathcal{H}_M}[\varphi], e' \rangle_{E, E'} = \sum_{n \in \mathbb{I}_+} \langle \phi_n, e' \rangle_{E, E'} \int_{\mathcal{S}} \langle \phi_n, \gamma s \rangle_{E, E'} \langle \varphi, \gamma s \rangle_{E, E'} d\nu(s).$$

Finally, because of the continuous inclusion of $\mathcal{H}_M, (\cdot| \cdot)_{\mathcal{H}}$ into $\overline{\mathcal{H}_M}^{\gamma, \nu}$, the considered sum also converges for $\|\cdot\|_{\gamma, \nu}$.

4.2. Truncated approach and approximation. — In practice, even if the spectral decomposition of the operator $\mathcal{L}_{\gamma, \nu} = \mathfrak{F}^t \mathfrak{F}$ is known, it is not always possible, for instance for numerical reasons, to consider all the terms appearing in the Mercer decomposition of $T_{\mathcal{H}_M}$,

$$\forall e' \in E', T_{\mathcal{H}_M} e' = \sum_{n \in \mathbb{I}_+} \lambda_n \langle \phi_n, e' \rangle_{E, E'} \phi_n.$$

A classic alternative simply consists in not considering all the terms that appear in the previous sum. In this case, one currently speaks about *spectral truncation* and these are usually the largest eigenvalues which are conserved. The following section aims at illustrating the use of this alternative in our context. Note that additional considerations concerning the optimal character of the approximation by truncation based on the largest eigenvalues are developed in Section 6.4.

Remark that we have to keep in mind that, in the most part of applications, the true analytical spectral decomposition of $\mathcal{L}_{\gamma, \nu}$ would be unknown. So, the study of the behavior of the proposed approach when dealing with approximated eigenpairs is of importance in regards of such applications.

4.2.1. *Spectral truncation.* — Assume that we dispose of an approximated kernel defined from a subset \mathbb{I}_{trc} of \mathbb{I}_+ , that is, for all $e' \in E'$,

$$T_{\mathcal{H}_M^{trc}} e' = \sum_{n \in \mathbb{I}_{trc}} \lambda_n \langle \phi_n, e' \rangle_{E, E'} \phi_n.$$

For $\varphi \in \mathcal{H}$, we then obtain an approximation of the optimal interpolator $P_{\mathcal{H}_M} [\varphi]$ that we denote by $P_{\mathcal{H}_M^{trc}} [\varphi]$. We have

$$\begin{aligned} \forall e' \in E', \langle P_{\mathcal{H}_M^{trc}} [\varphi], e' \rangle_{E, E'} &= \left(\varphi \middle| T_{\mathcal{H}_M^{trc}} e' \right)_{\mathcal{H}} \\ (4.2) \qquad \qquad \qquad &= \sum_{n \in \mathbb{I}_{trc}} \langle \phi_n, e' \rangle_{E, E'} \int_{\mathcal{S}} \tilde{\phi}_n(s) \langle \varphi, \gamma s \rangle_{E, E'} d\nu(s), \end{aligned}$$

where \mathcal{H}_M^{trc} is the closure in \mathcal{H} of the subspace spanned by the ϕ_n , for $n \in \mathbb{I}_{trc}$.

Theorem 4.2. — Consider the general framework of Section 4.2 and introduce the set $\mathbb{I}_{err} = \mathbb{I}_+ \setminus \mathbb{I}_{trc}$, then

$$(4.3) \qquad \forall e' \in E', \left\langle \mathbb{P}_{\mathcal{H}_M} [\varphi] - \mathbb{P}_{\mathcal{H}_M^{trc}} [\varphi], e' \right\rangle_{E, E'}^2 \leq \|\varphi\|_{\mathcal{H}}^2 \sum_{n \in \mathbb{I}_{err}} \lambda_n \langle \phi_n, e' \rangle_{E, E'}^2$$

$$(4.4) \qquad \text{and} \quad \left\| \varphi - \mathbb{P}_{\mathcal{H}_M^{trc}} [\varphi] \right\|_{\gamma, \nu}^2 \leq \|\varphi\|_{\mathcal{H}}^2 \sum_{n \in \mathbb{I}_{err}} \lambda_n.$$

Proof. — By definition, we have for all $e' \in E'$,

$$(4.5) \qquad \left\langle \mathbb{P}_{\mathcal{H}_M} [\varphi] - \mathbb{P}_{\mathcal{H}_M^{trc}} [\varphi], e' \right\rangle_{E, E'} = \left(\varphi \middle| \sum_{n \in \mathbb{I}_{err}} \lambda_n \langle \phi_n, e' \rangle_{E, E'} \phi_n \right)_{\mathcal{H}}.$$

To obtain expression (4.3), we just have to remark that the Cauchy-Schwarz inequality applied to equation (4.5) gives

$$\left\langle \mathbb{P}_{\mathcal{H}_M} [\varphi] - \mathbb{P}_{\mathcal{H}_M^{trc}} [\varphi], e' \right\rangle_{E, E'}^2 \leq \|\varphi\|_{\mathcal{H}}^2 \left\| \sum_{n \in \mathbb{I}_{err}} \lambda_n \langle \phi_n, e' \rangle_{E, E'} \phi_n \right\|_{\mathcal{H}}^2$$

and that, from Proposition 3.8,

$$\left\| \sum_{n \in \mathbb{I}_{err}} \lambda_n \langle \phi_n, e' \rangle_{E, E'} \phi_n \right\|_{\mathcal{H}}^2 = \sum_{n \in \mathbb{I}_{err}} \lambda_n \langle \phi_n, e' \rangle_{E, E'}^2.$$

Next, from equation (4.3) and the definition of $\|\cdot\|_{\gamma, \nu}^2$ (see section 3.3.2), we have,

$$\int_{\mathcal{S}} \left\langle \mathbb{P}_{\mathcal{H}_M} [\varphi] - \mathbb{P}_{\mathcal{H}_M^{trc}} [\varphi], \gamma s \right\rangle_{E, E'}^2 d\nu(s) \leq \|\varphi\|_{\mathcal{H}}^2 \sum_{n \in \mathbb{I}_{err}} \lambda_n \int_{\mathcal{S}} \langle \phi_n, \gamma s \rangle_{E, E'}^2 d\nu(s).$$

Since $\|\phi_n\|_{\gamma, \nu}^2 = 1$ for all $n \in \mathbb{I}_+$, we obtain

$$\left\| \mathbb{P}_{\mathcal{H}_M} [\varphi] - \mathbb{P}_{\mathcal{H}_M^{trc}} [\varphi] \right\|_{\gamma, \nu}^2 \leq \|\varphi\|_{\mathcal{H}}^2 \sum_{n \in \mathbb{I}_{err}} \lambda_n,$$

which concludes the proof. \square

Let us remark that, by definition of $\|\cdot\|_{\gamma,\nu}$,

$$(4.6) \quad \forall \varphi \in \mathcal{H}, \left\| \mathbb{P}_{\mathcal{H}_M}[\varphi] - \mathbb{P}_{\mathcal{H}_M^{trc}}[\varphi] \right\|_{\gamma,\nu}^2 = \left\| \varphi - \mathbb{P}_{\mathcal{H}_M^{trc}}[\varphi] \right\|_{\gamma,\nu}^2.$$

We call $\sum_{n \in \mathbb{I}_{err}} \lambda_n$ the *spectral error term*. It will be in practice evaluated by considering

$$(4.7) \quad \begin{aligned} \sum_{n \in \mathbb{I}_{err}} \lambda_n &= \sum_{n \in \mathbb{I}_+} \lambda_n - \sum_{n \in \mathbb{I}_{trc}} \lambda_n \\ &= \int_{\mathcal{S}} \|T_{\mathcal{H}} \gamma s\|_{\mathcal{H}}^2 d\nu(s) - \sum_{n \in \mathbb{I}_{trc}} \lambda_n. \end{aligned}$$

A good indicator (see also Section 6.4) of the overall quality of the obtained approximation can classically be found in the ratios

$$\frac{\sum_{n \in \mathbb{I}_{trc}} \lambda_n}{\sum_{n \in \mathbb{I}_+} \lambda_n} = 1 - \frac{\sum_{n \in \mathbb{I}_{err}} \lambda_n}{\sum_{n \in \mathbb{I}_+} \lambda_n}.$$

4.3. About the choice of the parameterization. — In this section, we mention some general considerations concerning the choice of the parameterization. By parameterization, we mean here the overall process leading to the construction of a regular embedding of \mathcal{H}_M into an auxiliary Hilbert space.

4.3.1. Computational aspects. — In the interpolation context of Theorem 4.1, the parameterization can just appear as a tool allowing to obtain the representation formula (4.1). No matter its choice from the moment it allows the definition of a regular embedding \mathfrak{F} of \mathcal{H}_M .

Nevertheless, if one envisages the computation of the elements constituting the orthonormal basis of \mathcal{H}_M associated with the considered embedding (Proposition 3.8), this choice takes importance. Indeed, it in part determines the operator which has to be diagonalized. It then appears reasonable to try to make a choice that defines a simplest as possible spectral problem. One will for instance try to be in a position allowing an analytical resolution, or the use of a particular numerical method.

In such a context, an illustration of what appears to us as relatively judicious choices of parameterizations can be found in [Gau11, Section 3.3]. In this particular example, appropriate choices allow to obtain analytical expressions for many of the involved quantities and prediction formulae concerning the conditioning of a two parameters Brownian sheet are, in so doing, obtained in an original way.

4.3.2. The approximation case. — In addition of this first consideration, the parametrization choice has a direct influence on the behavior of the optimal interpolator approximation obtained by spectral truncation in equation (4.2). Indeed, different choices of parameterization for a same problem lead to different approximations of the optimal interpolator, and this even if the spectral ratios of the considered truncations are equal.

The parameterization directly influences the way $\mathbb{P}_{\mathcal{H}_M^{trc}}[\varphi]$ approximates $\mathbb{P}_{\mathcal{H}_M}[\varphi]$ on M . So, it offers a way to modulate the accuracy of the approximation in function

of the elements of M . This characteristic of *tunable precision* could offer interesting possibilities in regards of applications.

5. Finite Case

5.1. Context and notations. — We suppose that M is of finite dimension, *i.e.* $M = \text{span}\{\mu_1, \dots, \mu_n\}$, with $n \in \mathbb{N}^*$. Let us define the matrix $\mathbf{T} \in \mathbb{R}^{n \times n}$ by,

$$(5.1) \quad \text{for } 1 \leq i, j \leq n, \mathbf{T}_{i,j} = (T_{\mathcal{H}}\mu_i | T_{\mathcal{H}}\mu_j)_{\mathcal{H}}.$$

For simplicity and without loss of generality, we assume that the $\mu_i \in E'$ are such that the symmetric and positive matrix \mathbf{T} is *invertible*. For convenience, we introduce the following matrix notations,

$$\boldsymbol{\mu} = (\mu_1, \dots, \mu_n)^T \text{ and } \mathbf{T} = (T_{\mathcal{H}}\boldsymbol{\mu} | T_{\mathcal{H}}\boldsymbol{\mu}^T)_{\mathcal{H}} = \langle T_{\mathcal{H}}\boldsymbol{\mu}, \boldsymbol{\mu}^T \rangle = \langle \boldsymbol{\mu}, T_{\mathcal{H}}\boldsymbol{\mu}^T \rangle,$$

where $T_{\mathcal{H}}\boldsymbol{\mu} = (T_{\mathcal{H}}\mu_1, \dots, T_{\mathcal{H}}\mu_n)^T$ is a column vector. Hence, for $\varphi \in \mathcal{H}$, the optimal interpolator of φ for M can be written under the form

$$(5.2) \quad h_{\varphi, M} = T_{\mathcal{H}}\boldsymbol{\mu}^T \mathbf{T}^{-1} \langle \boldsymbol{\mu}, \varphi \rangle,$$

with $\langle \boldsymbol{\mu}, \varphi \rangle = \left(\langle \varphi, \mu_1 \rangle_{E, E'}, \dots, \langle \varphi, \mu_n \rangle_{E, E'} \right)^T$. Remark for instance that with our notations, $\langle \boldsymbol{\mu}, \varphi \rangle^T = \langle \varphi, \boldsymbol{\mu}^T \rangle$ and, for $e \in E$ and $e' \in E'$, $\langle e, e' \rangle_{E, E'} = \langle e, e' \rangle = \langle e', e \rangle$. So, we will write, for $e' \in E'$, $\langle h_{\varphi, M}, e' \rangle_{E, E'} = \langle e', T_{\mathcal{H}}\boldsymbol{\mu}^T \rangle \mathbf{T}^{-1} \langle \boldsymbol{\mu}, \varphi \rangle$.

The aim of this section is to prove, by explicit calculations, that the expression of the minimal norm interpolator given in equation (5.2) is equal to the one given in equation (4.1) of Theorem 4.1.

5.2. Parameterization. — We first introduce a trivial parameterization of this problem. Let $\mathcal{S} = \{1, \dots, n\}$ and consider the measure ν on \mathcal{S} which assigns a weight $w_i > 0$ to each $i \in \mathcal{S} = \{1, \dots, n\}$. The auxiliary space $L^2(\mathcal{S}, \nu)$ can then be identified to the space \mathbb{R}^n endowed with the inner-product $(\mathbf{x} | \mathbf{y})_{\mathbf{W}} = \mathbf{x}^T \mathbf{W} \mathbf{y}$, where \mathbf{x} and \mathbf{y} are two column vectors of \mathbb{R}^n and where \mathbf{W} is the matrix

$$\mathbf{W} = \text{diag}(w_1, \dots, w_n).$$

Notice that we identify a vector of \mathbb{R}^n with the column vector of its coefficients in the canonical basis of \mathbb{R}^n .

We next consider the application $\gamma : \mathcal{S} \rightarrow E'$, given by $\gamma i = \mu_i$, for all $i \in \{1, \dots, n\}$. The associated application \mathfrak{F} is trivially a regular embedding associated with our problem. It verifies, for all $h \in \mathcal{H}$, $\mathfrak{F}h = \langle \boldsymbol{\mu}, h \rangle \in \mathbb{R}^n$ (and $\mathfrak{F}h(i) = \langle h, \mu_i \rangle_{E, E'}$, for all $i \in \{1, \dots, n\} = \mathcal{S}$). If we identify $L^2(\mathcal{S}, \nu)$ with \mathbb{R}^n , then for $\boldsymbol{\alpha} \in \mathbb{R}^n$,

$$(5.3) \quad {}^t \mathfrak{F} \boldsymbol{\alpha} = T_{\mathcal{H}}\boldsymbol{\mu}^T \mathbf{W} \boldsymbol{\alpha} \in \mathcal{H}_M.$$

We finally obtain that $\mathcal{L}_{\gamma, \nu} = \mathfrak{F}^t \mathfrak{F}$ is given by

$$(5.4) \quad \mathfrak{F}^t \mathfrak{F} \boldsymbol{\alpha} = \mathbf{T} \mathbf{W} \boldsymbol{\alpha}.$$

In the same way, we have (see Proposition 3.9),

$$\begin{aligned}
 \forall h \in \mathcal{H}, L_{\gamma, \nu}[h] &= {}^t \mathfrak{F} \mathfrak{F} h = \int_S \langle h, \gamma s \rangle_{E, E'} T_{\mathcal{H}} \gamma s \, d\nu(s) \\
 (5.5) \qquad \qquad \qquad &= \sum_{i=1}^n w_i \langle h, \mu_i \rangle_{E, E'} T_{\mathcal{H}} \mu_i = T_{\mathcal{H}} \boldsymbol{\mu}^T \mathbf{W} \langle \boldsymbol{\mu}, h \rangle.
 \end{aligned}$$

The symmetric and positive bilinear form $(\cdot | \cdot)_{\gamma, \nu}$ on \mathcal{H} , associated with \mathfrak{F} via equation (3.18), is given by, for h and $g \in \mathcal{H}$,

$$(h | g)_{\gamma, \nu} = \sum_{i=1}^n w_i \langle h, \mu_i \rangle_{E, E'} \langle g, \mu_i \rangle_{E, E'} = \langle h, \boldsymbol{\mu}^T \rangle \mathbf{W} \langle \boldsymbol{\mu}, g \rangle.$$

5.3. Spectral decomposition. — Let $\lambda_1 > 0, \dots, \lambda_n > 0$ be the eigenvalues of $\mathbf{T}\mathbf{W}$ and let $\mathbf{v}_1, \dots, \mathbf{v}_n$ be their associated eigenvectors, *i.e.* $\mathbf{T}\mathbf{W} = \mathbf{P}\boldsymbol{\Lambda}\mathbf{P}^{-1}$ with

$$\boldsymbol{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_n) \text{ and } \mathbf{P} = (\mathbf{v}_1 | \dots | \mathbf{v}_n).$$

Note that $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ forms an orthonormal basis of \mathbb{R}^n , $(\cdot | \cdot)_{\mathbf{W}}$, *i.e.*

$$(5.6) \qquad \mathbf{P}^T \mathbf{W} \mathbf{P} = \text{Id}_{n \times n}, \text{ where } \text{Id}_{n \times n} \text{ is the } n \times n \text{ identity matrix.}$$

For $k \in \{1, \dots, n\}$, we define

$$(5.7) \qquad \phi_k = \frac{1}{\lambda_k} {}^t \mathfrak{F} \mathbf{v}_k = \frac{1}{\lambda_k} T_{\mathcal{H}} \boldsymbol{\mu}^T \mathbf{W} \mathbf{v}_k \in \mathcal{H}_M.$$

Proposition 5.1. — *Under the assumptions of Section 5, we have, for all $\varphi \in \mathcal{H}$,*

$$T_{\mathcal{H}} \boldsymbol{\mu}^T \mathbf{T}^{-1} \langle \boldsymbol{\mu}, \varphi \rangle = \sum_{k=1}^n \phi_k \int_S \langle \phi_k, \gamma s \rangle_{E, E'} \langle \varphi, \gamma s \rangle_{E, E'} \, d\nu(s).$$

Proof. — We have

$$\begin{aligned}
 T_{\mathcal{H}} \boldsymbol{\mu}^T \mathbf{T}^{-1} \langle \boldsymbol{\mu}, \varphi \rangle &= T_{\mathcal{H}} \boldsymbol{\mu}^T \mathbf{W} \mathbf{W}^{-1} \mathbf{T}^{-1} \langle \boldsymbol{\mu}, \varphi \rangle \\
 &= T_{\mathcal{H}} \boldsymbol{\mu}^T \mathbf{W} \mathbf{P} \boldsymbol{\Lambda}^{-1} \mathbf{P}^{-1} \langle \boldsymbol{\mu}, \varphi \rangle.
 \end{aligned}$$

Let us study the terms appearing in this last expression. First, from equation (5.7),

$$T_{\mathcal{H}} \boldsymbol{\mu}^T \mathbf{W} \mathbf{P} \boldsymbol{\Lambda}^{-1} = (\phi_1, \dots, \phi_n).$$

Next, using equation (5.6), we find

$$\mathbf{P}^{-1} \langle \boldsymbol{\mu}, \varphi \rangle = (\mathbf{P}^T \mathbf{W} \mathbf{P})^{-1} \mathbf{P}^T \mathbf{W} \langle \boldsymbol{\mu}, \varphi \rangle = \mathbf{P}^T \mathbf{W} \langle \boldsymbol{\mu}, \varphi \rangle.$$

To conclude, we remark that, for $k \in \{1, \dots, n\}$, the term

$$\int_S \langle \phi_k, \gamma s \rangle_{E, E'} \langle \varphi, \gamma s \rangle_{E, E'} \, d\nu(s) = \sum_{i=1}^n w_i \langle \varphi, \mu_i \rangle_{E, E'} \langle \phi_k, \mu_i \rangle_{E, E'}$$

is the k -th component of the vector $\mathbf{P}^T \mathbf{W} \langle \boldsymbol{\mu}, \varphi \rangle$. □

6. Application to Gaussian process models

Optimal interpolation in Hilbert subspaces and Gaussian process models are intrinsically linked. In this section, we recall some of the main properties concerning the conditioning of Gaussian processes in the framework of topological vector spaces. We also apply the spectral approach developed in Sections 3 and 4 to the conditioning problem. The IMSE-optimal character of the approximation by truncation is finally addressed in Section 6.4.

6.1. Notations and recalls. — Let \mathcal{H} be a *separable* Hilbert subspace of E . Throughout Section 6, we assume that \mathcal{H} is the Cameron-Martin space of a centered Gaussian process Y defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. We denote by \mathbf{H} the Gaussian Hilbert space associated with Y (see [Jan97]). We remind that \mathbf{H} is a closed linear subspace of the Hilbert space $L^2(\Omega, \mathcal{F}, \mathbb{P})$ of second order real random variables (r.v.) on $(\Omega, \mathcal{F}, \mathbb{P})$ (we identify random variables that are equal \mathbb{P} -almost surely).

For the sake of simplicity, we assume that E is a Banach space (see [DFLC71, Tal83] for more general frameworks). We also consider that Y takes its values in E with \mathbb{P} -probability 1 (*i.e.* the triplet $(j, \mathcal{H}, \overline{\mathcal{H}}^E)$ is an *abstract Wiener space*, where $\overline{\mathcal{H}}^E$ is the completed of \mathcal{H} in E and where j is the continuous inclusion of \mathcal{H} into $\overline{\mathcal{H}}^E$, see [Gro67]).

We denote by $\mathcal{I} : \mathcal{H} \rightarrow \mathbf{H}$ the isometry between \mathcal{H} and \mathbf{H} , it verifies

$$\mathbb{E}(\mathcal{I}h\mathcal{I}g) = (h|g)_{\mathcal{H}},$$

where $\mathbb{E}(\mathcal{I}h\mathcal{I}g)$ represents the inner-product in $L^2(\Omega, \mathcal{F}, \mathbb{P})$ between the two centered random variables $\mathcal{I}h$ and $\mathcal{I}g \in \mathbf{H}$. Let us also add, for $e' \in E'$,

$$\langle Y, e' \rangle_{E, E'} \stackrel{(\text{notation})}{=} Y_{e'} = \mathcal{I}(T_{\mathcal{H}}e').$$

One can consult, among others, [Bax76, TV07, Gau11] for more details about the previous notions.

For a linear subspace M of E' , $P_{\mathcal{H}_M}$ denotes the orthogonal projection of \mathcal{H} onto \mathcal{H}_M . We then introduce the orthogonal projection $P_{\mathbf{H}_M}$ of \mathbf{H} onto $\mathbf{H}_M = \mathcal{I}(\mathcal{H}_M)$. We have the commutative diagram

$$(6.1) \quad \begin{array}{ccc} \mathcal{H} & \xrightarrow{\mathcal{I}} & \mathbf{H} \\ \downarrow P_{\mathcal{H}_M} & & \downarrow P_{\mathbf{H}_M} \\ \mathcal{H}_M & \xrightarrow{\mathcal{I}} & \mathbf{H}_M \end{array}$$

The application $T_{\mathcal{H}_M} = P_{\mathcal{H}_M}T_{\mathcal{H}}$ is the Hilbert kernel of \mathcal{H}_M , hence, by isometry,

$$(6.2) \quad \forall e' \in E', \mathcal{I}(T_{\mathcal{H}_M}e') = P_{\mathbf{H}_M}[Y_{e'}] \stackrel{(\text{notation})}{=} \mathbb{E}(Y_{e'}|Y_{f'}, f' \in M).$$

For all $e' \in E'$, the r.v. $P_{\mathbf{H}_M}[Y_{e'}]$ is called the *conditional mean of $Y_{e'}$ knowing $Y_{f'}$ for $f' \in M$* and $T_{\mathcal{H}_0}$ is the associated *conditional covariance kernel*. The notion of *conditional law* of the process Y is addressed in Section 6.3.

6.2. Spectral approach for conditioning. — We now consider the general framework of Section 3.

Proposition 6.1. — *Let us consider the centered Gaussian process $(Y_{\gamma s})_{s \in \mathcal{S}}$. Under the assumptions of Theorem 4.1, the sample paths of $(Y_{\gamma s})_{s \in \mathcal{S}}$ are in $L^2(\mathcal{S}, \nu)$ with \mathbb{P} -probability 1. In addition,*

$$\mathbb{E} \left[\int_{\mathcal{S}} (Y_{\gamma s})^2 d\nu(s) \right] = \sum_{n \in \mathbb{I}_+} \lambda_n (= \tau).$$

Proof. — For $s \in \mathcal{S}$, we remind that $Y_{\gamma s} = \mathcal{I}(T_{\mathcal{H}} \gamma s)$. Let $\{h_j : j \in \mathbb{J}\}$ be an orthonormal basis of \mathcal{H} . As in equation (2.2), $(Y_{\gamma s})_{s \in \mathcal{S}}$ admits the Karhunen-Loève expansion

$$(6.3) \quad \forall s \in \mathcal{S}, Y_{\gamma s} = \sum_{j \in \mathbb{J}} \langle h_j, \gamma s \rangle_{E, E'} \mathcal{I}(h_j),$$

where the $\mathcal{I}(h_j) = \zeta_j$, $j \in \mathbb{J}$, form by isometry an orthonormal basis of \mathbf{H} (such independent $\mathcal{N}(0, 1)$ r.v. are sometime called *orthogaussian*, see [Dud10]). We then deduce from Condition C-i that the sample paths of $(Y_{\gamma s})_{s \in \mathcal{S}}$ are measurable (as real-valued functions on \mathcal{S}) with \mathbb{P} -probability 1.

One can obviously choose the orthonormal basis $\{h_j : j \in \mathbb{J}\}$ of \mathcal{H} such that it coincides on \mathcal{H}_M with the orthonormal basis $\{\sqrt{\lambda_n} \phi_n : n \in \mathbb{I}_+\}$ associated with the considered regular embedding \mathfrak{F} (Proposition 3.8). Then, from C-iii and C-iv,

$$\begin{aligned} \sum_{j \in \mathbb{J}} \int_{\mathcal{S}} \mathbb{E} \left[(\mathfrak{F} h_j(s) \zeta_j)^2 \right] d\nu(s) &= \sum_{j \in \mathbb{J}} \|\mathfrak{F} h_j\|_{L^2}^2 \\ &= \sum_{n \in \mathbb{I}_+} \left\| \sqrt{\lambda_n} \tilde{\phi}_n \right\|_{L^2}^2 = \sum_{n \in \mathbb{I}_+} \lambda_n < +\infty. \end{aligned}$$

Thus, the sum $\sum_{j \in \mathbb{J}} \|\mathfrak{F} h_j \zeta_j\|_{L^2(\nu \otimes \mathbb{P})}^2$ is convergent, which, from Tonelli's Theorem, implies the convergence in $L^2(\nu \otimes \mathbb{P})$ of

$$(s, \omega) \mapsto \sum_{j \in \mathbb{J}} \mathfrak{F} h_j(s) \zeta_j(\omega), \text{ with } s \in \mathcal{S} \text{ and } \omega \in \Omega$$

and finally completes the proof. \square

Theorem 6.2. — *Let Y be a centered Gaussian process with values in E and covariance kernel $T_{\mathcal{H}}$. Let M be a linear subspace of E' . Under the assumptions of Theorem 4.1, we have, for all $e' \in E'$,*

$$(6.4) \quad \mathbb{E}(Y_{e'} | Y_{f'}, f' \in M) = \sum_{n \in \mathbb{I}_+} \langle \phi_n, e' \rangle_{E, E'} \int_{\mathcal{S}} \tilde{\phi}_n(s) Y_{\gamma s} d\nu(s).$$

In addition, the centered Gaussian process with covariance kernel $T_{\mathcal{H}_M}$ (that is the process corresponding to $\mathbb{E}(Y_{e'} | Y_{f'}, f' \in M)$, with $e' \in E'$) takes its values in $\overline{\mathcal{H}_M}^{\gamma, \nu}$ with \mathbb{P} -probability 1.

Proof. — We have to verify that the right member of equation (6.4) is well-defined. From Proposition 3.8, we know that, for all $e' \in E'$,

$$T_{\mathcal{H}_M} e' = \sum_{n \in \mathbb{I}_+} \lambda_n \langle \phi_n, e' \rangle_{E, E'} \phi_n,$$

this series being convergent in \mathcal{H} , so, by isometry,

$$(6.5) \quad \mathcal{I}(T_{\mathcal{H}_M} e') = \sum_{n \in \mathbb{I}_+} \lambda_n \langle \phi_n, e' \rangle_{E, E'} \mathcal{I}(\phi_n).$$

For $n \in \mathbb{I}_+$, we have, thanks to equation (3.12),

$$\phi_n = \frac{1}{\lambda_n} \int_{\mathcal{S}} \tilde{\phi}_n(s) T_{\mathcal{H}} \gamma_s d\nu(s).$$

Now, Proposition 6.1 assures that the expression

$$(6.6) \quad \mathcal{I}(\phi_n) = \frac{1}{\lambda_n} \int_{\mathcal{S}} \tilde{\phi}_n(s) \mathcal{I}(T_{\mathcal{H}} \gamma_s) d\nu(s) = \frac{1}{\lambda_n} \int_{\mathcal{S}} \tilde{\phi}_n(s) Y_{\gamma_s} d\nu(s)$$

keeps sense under our working hypotheses. At last, from expansion (6.4), Proposition 6.1 and the isometry between $\overline{\mathfrak{F}(\mathcal{H})}^{L^2}$ and $\overline{\mathcal{H}_M}^{\gamma, \nu}$, we deduce that the process $\mathbb{E}(Y_{e'} | Y_{f'}, f' \in M)$ takes its values in $\overline{\mathcal{H}_M}^{\gamma, \nu}$ with \mathbb{P} -probability 1. \square

6.3. A note on regular conditional probabilities. — We now study the conditional laws of the process Y relatively to the knowledge of the values take by its sample paths on M . Because, in our study, \mathcal{H}_M can be infinite dimensional, the construction of a regular conditional probability requires some precautions. We consider a case where such a conditional probability exists (see for instance [TV07], this is for instance always true if \mathcal{H}_M is finite dimensional). We finally introduce an additional assumption which assures the existence of a spectral representation for the mean of the conditional laws, this representation corresponding to a natural extension of the one obtained in equation (4.1).

Let us assume that

$$(6.7) \quad \overline{\mathcal{H}}^E = \overline{\mathcal{H}_M}^E \oplus \overline{\mathcal{H}_0}^E,$$

which means that $\overline{\mathcal{H}}^E = \overline{\mathcal{H}_M}^E + \overline{\mathcal{H}_0}^E$ with $\overline{\mathcal{H}_M}^E \cap \overline{\mathcal{H}_0}^E = \{0\}$.

Condition (6.7) assures the existence of the linear continuous projection \mathfrak{P} of $\overline{\mathcal{H}}^E$ onto $\overline{\mathcal{H}_M}^E$ parallel to $\overline{\mathcal{H}_0}^E$ (i.e. $\mathfrak{P}h_0 = 0$ for all $h_0 \in \overline{\mathcal{H}_0}^E$).

We then consider the family of Gaussian measures on $\overline{\mathcal{H}}^E$ with mean $\mathfrak{P}[\Psi] \in \overline{\mathcal{H}_M}^E$, for $\Psi \in \overline{\mathcal{H}}^E$ and covariance kernel $T_{\mathcal{H}_0}$. From [TV07, Theorem 3.11], such a family defines a *regular conditional probability* over $\overline{\mathcal{H}}^E$ relative to the knowledge of Y on M . In such case, the following notation is often used:

$$(6.8) \quad \mathfrak{P}[\Psi] = \mathbb{E}(Y_{e'} | Y_{f'} = \langle \Psi, f' \rangle_{E, E'}, f' \in M).$$

Note that if we denote by μ_Y the Gaussian measure on $\overline{\mathcal{H}}^E$ associated with Y , the preceding regular conditional probability corresponds to the *disintegration* of μ_Y relatively to \mathfrak{P} .

Now, let us also suppose that $\overline{\mathcal{H}_M}^E$ can be continuously injected into $\overline{\mathcal{H}_M}^{\gamma, \nu}$, what we write

$$(6.9) \quad \overline{\mathcal{H}_M}^E \hookrightarrow \overline{\mathcal{H}_M}^{\gamma, \nu}.$$

We next consider the extension $\tilde{\mathfrak{F}}$ of \mathfrak{F} to $\overline{\mathcal{H}}^E$ defines by $\tilde{\mathfrak{F}}h_0 = 0$, for all $h_0 \in \overline{\mathcal{H}_0}^E$ and, on $\overline{\mathcal{H}_M}^E$, by the continuous extension of the restriction of \mathfrak{F} to \mathcal{H}_M . From Section 3.3.3 and condition (6.9), $\tilde{\mathfrak{F}}$ is well-defined from $\overline{\mathcal{H}}^E$ onto $\overline{\mathfrak{F}(\mathcal{H})}^{L^2}$.

Then, as for $P_{\mathcal{H}_M} = \mathfrak{F}^\dagger \mathfrak{F}$, we have $\mathfrak{P} = \mathfrak{F}^\dagger \tilde{\mathfrak{F}}$ (remark that this last expression is well-defined in regards of the definition of \mathfrak{F}^\dagger given in Section 3.3.4). We finally obtain the spectral representation formula,

$$(6.10) \quad \forall \Psi \in \overline{\mathcal{H}}^E, \mathfrak{P}[\Psi] = \sum_{n \in \mathbb{I}_+} \langle \phi_n, e' \rangle_{E, E'} \int_{\mathcal{S}} \tilde{\phi}_n(s) \langle \Psi, \gamma s \rangle_{E, E'} d\nu(s).$$

Remark 6.3. — We have the well-known equality, for $\varphi \in \mathcal{H}$ and $e' \in E'$,

$$(6.11) \quad \langle P_{\mathcal{H}_M}[\varphi], e' \rangle_{E, E'} = \mathbb{E}(Y_{e'} | Y_{f'} = \langle \varphi, f' \rangle_{E, E'}, f' \in M).$$

6.4. Optimal approximation. — The results and considerations of Section 4.2 can also be extended to the Gaussian processes case. In this section, we discuss the optimal character of the approximation by truncation.

Let Y be a centered Gaussian process with values in E and covariance $T_{\mathcal{H}}$. Under the assumptions of Theorem 6.2, we consider a subset \mathbb{I}_{trc} of \mathbb{I}_+ composed of the largest eigenvalues of $\mathcal{L}_{\gamma, \nu}$, in the sense that,

$$(6.12) \quad \text{if } i \in \mathbb{I}_{err} = \mathbb{I}_+ \setminus \mathbb{I}_{trc} \text{ and } n \in \mathbb{I}_{trc}, \text{ then } \lambda_i \leq \lambda_n.$$

Proposition 6.4. — Let \mathcal{H}_M^{app} be any closed linear subspace of \mathcal{H}_M and denote by $T_{\mathcal{H}_M^{app}}$ the associated Hilbert kernel. For $e' \in E'$, we introduce the two approximations of $Z_{e'} = \mathbb{E}(Y_{e'} | Y_{f'} = \langle \varphi, f' \rangle_{E, E'}, f' \in M)$,

$$(6.13) \quad Z_{e'}^{trc} = \mathcal{I}(T_{\mathcal{H}_M^{trc}} e') \text{ and } Z_{e'}^{app} = \mathcal{I}(T_{\mathcal{H}_M^{app}} e').$$

If \mathcal{H}_M^{trc} and \mathcal{H}_M^{app} have the same finite dimension $N \in \mathbb{N}^*$, then under equation (6.12),

$$(6.14) \quad \mathbb{E}(\|Y - Z^{trc}\|_{\gamma, \nu}^2) \leq \mathbb{E}(\|Y - Z^{app}\|_{\gamma, \nu}^2).$$

Proof. — From Theorem 4.2 and Proposition 6.1, we have

$$(6.15) \quad \mathbb{E}(\|Z^{trc}\|_{\gamma, \nu}^2) = \mathbb{E}\left(\int_{\mathcal{S}} (Z_{\gamma s}^{trc})^2 d\nu(s)\right) = \int_{\mathcal{S}} \|T_{\mathcal{H}_M^{trc}} \gamma s\|_{\mathcal{H}}^2 d\nu(s) = \sum_{n \in \mathbb{I}_{trc}} \lambda_n$$

$$\text{and } \mathbb{E}(\|Y - Z^{trc}\|_{\gamma, \nu}^2) = \mathbb{E}(\|Y\|_{\gamma, \nu}^2) - \mathbb{E}(\|Z^{trc}\|_{\gamma, \nu}^2).$$

Let f_1, \dots, f_N be an orthonormal basis of \mathcal{H}_M^{app} and consider its decomposition in the orthonormal basis associated with the considered regular embedding, that is

$$\forall i \in \{1, \dots, N\}, f_i = \sum_{k \in \mathbb{I}_+} \alpha_{i,k} \sqrt{\lambda_k} \phi_k, \text{ with } \alpha_{i,k} \in \mathbb{R}$$

and where, for i and $j \in \{1, \dots, N\}$, $\sum_{k \in \mathbb{I}_+} \alpha_{i,k} \alpha_{j,k} = \delta_{i,j}$ (Kronecker delta). Thus, we easily obtain that

$$\mathbb{E}(\|Z^{app}\|_{\gamma,\nu}^2) = \sum_{i=1}^N \sum_{k \in \mathbb{I}_+} \alpha_{i,k}^2 \lambda_k.$$

Next, using for instance convex combinations arguments, we remark that

$$(6.16) \quad \sum_{n \in \mathbb{I}_{trc}} \lambda_n \geq \sum_{i=1}^N \sum_{k \in \mathbb{I}_+} \alpha_{i,k}^2 \lambda_k.$$

We finally conclude thanks to equation (6.15). \square

In the same way as in equation (4.6), one can remark that

$$(6.17) \quad \mathbb{E}(\|Y - Z^{app}\|_{\gamma,\nu}^2) = \mathbb{E}(\|Z - Z^{app}\|_{\gamma,\nu}^2).$$

Thus, in regards of $\|\cdot\|_{\gamma,\nu}$ (*i.e.* in the sense of equation (6.14)), for all $e' \in E'$, $Z_{e'}^{trc}$ is the best approximation, based on N elements of \mathcal{H}_M , of the conditional mean $\mathbb{E}(Y_{e'} | Y_{f'} = \langle \varphi, f' \rangle_{E,E'}, f' \in M)$. One can hence speak about a certain *IMSE-optimality* (Integrated Mean Square Error) of the approximation by truncation. This point is illustrated in Section 7.5.3.

7. Example of application

7.1. The problem. — Let $\mathcal{X} = \mathbb{R}^2$ and \mathcal{H} be the RKHS of real-valued functions on \mathcal{X} (see Section 2.1.2) associated with the kernel (squared exponential or Gaussian kernel, see for instance [RW06]), for x and $y \in \mathcal{X}$,

$$K(x, y) = e^{-\frac{\|x-y\|^2}{\sigma^2}}, \text{ with } \sigma > 0 \text{ and } \|\cdot\| \text{ the euclidean norm.}$$

For $m \in \mathbb{N}$, let $\mathcal{E}^m \subset \mathbb{R}^{\mathcal{X}}$ be the subspace of functions of class C^m endowed with the topology of the uniform convergence on the compact subsets of \mathcal{X} for all the derivatives of order $\leq m$ (of general order if $m = +\infty$). From [Sch64, Proposition 25], for all $m \in \mathbb{N}$ (and also for $m = +\infty$), \mathcal{H} is a Hilbert subspace of \mathcal{E}^m . In what follows, we will consider \mathcal{H} as a Hilbert subspace of $E = \mathcal{E}^1$.

Let $x = (x_1, x_2) \in \mathcal{X}$, we will use polar coordinates, *i.e.* $x = (r_x \cos \alpha_x, r_x \sin \alpha_x)$ with $r_x \in \mathbb{R}_+$ and $\alpha_x \in [0, 2\pi)$. For $x \in \mathcal{X}$, we define $\delta_x \in E'$ and $\eta_x \in E'$ by

$$\forall f \in E, \langle f, \delta_x \rangle_{E,E'} = f(x) \text{ and } \langle f, \eta_x \rangle_{E,E'} = \frac{\partial f}{\partial r_x}(x),$$

δ_x is the Dirac measure centered on x and η_x corresponds to the evaluation of the radial derivative at x .

Let $\mathcal{C} \subset \mathbb{R}^2$ be the circle of center 0 and radius $R > 0$. We consider the linear subspaces of E'

$$M_D = \text{span}\{\delta_t, t \in \mathcal{C}\} \text{ and } M_N = \text{span}\{\eta_t, t \in \mathcal{C}\}$$

and $M_C = M_D + M_N$ (D and N stand for *Dirichlet* and *Neumann conditions*, C for *Cauchy*). The aim of this example is to approximate the kernel $K_{0_C}(\cdot, \cdot)$ of the subspace \mathcal{H}_{0_C} of functions $h \in \mathcal{H}$ such that,

$$\forall t \in \mathcal{C}, \langle h, \delta_t \rangle_{E, E'} = 0 \text{ and } \langle h, \eta_t \rangle_{E, E'} = 0.$$

Nevertheless, let us remark that what follows contains all the necessary informations to tackle the general interpolation problem associated with \mathcal{H} , an element $\varphi \in \mathcal{H}$ and M_C (see [Gau11]).

We present a two steps methodology. The first step (Section 7.2) consists in considering independently the interpolation problems in \mathcal{H} associated with M_N and M_D . Thanks to the study of a third operator (Section 7.3), we finally combine our results in Section 7.4 and obtain a model in which both values and radial derivatives are controlled on the circle (*Cauchy condition*). Numerical computations are finally presented in Section 7.5.

7.2. The two independent problems. — Let us introduce the linear subspaces of \mathcal{H} naturally associated with M_D and M_N ,

$$\begin{aligned} \mathcal{H}_{M_D} &= \mathcal{H}_{0_D}^\perp = \overline{\text{span} \{K(t, \cdot), t \in \mathcal{C}\}}^{\mathcal{H}} \text{ and} \\ \mathcal{H}_{M_N} &= \mathcal{H}_{0_N}^\perp = \overline{\text{span} \left\{ \frac{\partial K}{\partial r_t}(t, \cdot), t \in \mathcal{C} \right\}}^{\mathcal{H}}. \end{aligned}$$

We denote by $T_{\mathcal{H}_{M_D}}$ and $K_{M_D}(\cdot, \cdot)$ the Hilbert kernel and the reproducing kernel of \mathcal{H}_{M_D} respectively. We use similar notations for the kernels associated with M_N .

7.2.1. Parameterization. — Let $\mathcal{S} = [0, 2\pi)$ endowed with its natural Lebesgue measure (up to the multiplicative constant R) and consider the Hilbert space $L^2([0, 2\pi))$ of square-integrable real-valued functions (with respect to the Lebesgue measure) on $[0, 2\pi)$, endowed with the norm

$$\forall f \in L^2([0, 2\pi)), \|f\|_{L^2}^2 = \int_0^{2\pi} f(\theta)^2 R d\theta,$$

$L^2([0, 2\pi))$ plays the role of the auxiliary Hilbert space defined in Section 3.

We pose $s_{R, \theta} = (R \cos \theta, R \sin \theta) \in \mathcal{C}$ and introduce

$$\gamma_D : [0, 2\pi) \rightarrow M_D, \theta \mapsto \delta_{s_{R, \theta}} \text{ and } \gamma_N : [0, 2\pi) \rightarrow M_N, \theta \mapsto \eta_{s_{R, \theta}}.$$

We easily verify that γ_D and γ_N define two regular embeddings \mathfrak{F}_D and \mathfrak{F}_N of \mathcal{H}_{M_D} and \mathcal{H}_{M_N} into $L^2([0, 2\pi))$ (see also Remark 7.2) and which are given by, for $h \in \mathcal{H}$ and $\theta \in [0, 2\pi)$,

$$\mathfrak{F}_D h(\theta) = \langle h, \gamma_D \theta \rangle_{E, E'} \text{ and } \mathfrak{F}_N h(\theta) = \langle h, \gamma_N \theta \rangle_{E, E'}.$$

The associated integral operators $\mathcal{L}_D = \mathfrak{F}_D^t \mathfrak{F}_D$ and $\mathcal{L}_N = \mathfrak{F}_N^t \mathfrak{F}_N$ are defined on $L^2([0, 2\pi))$ and

$$(7.1) \quad \mathcal{L}_D[f](\alpha) = \int_0^{2\pi} K(x_{R, \alpha}, s_{R, \theta}) f(\theta) R d\theta,$$

$$(7.2) \quad \mathcal{L}_N[f](\alpha) = \int_0^{2\pi} \frac{\partial^2 K}{\partial r_s \partial r_x}(x_{R,\alpha}, s_{R,\theta}) f(\theta) R d\theta,$$

where $x_{R,\alpha} = (R \cos \alpha, R \sin \alpha) \in \mathcal{C}$, $\alpha \in [0, 2\pi)$ and $f \in L^2([0, 2\pi))$.

Remark 7.1. — Let $x = (r_x \cos \alpha_x, r_x \sin \alpha_x)$ and $y = (r_y \cos \alpha_y, r_y \sin \alpha_y)$ be two points of \mathcal{X} , we have

$$K(x, y) = \exp \left\{ -\frac{1}{\sigma^2} (r_x^2 + r_y^2 - 2r_x r_y \cos(\alpha_x - \alpha_y)) \right\},$$

$$\frac{\partial K}{\partial r_x}(x, y) = -\frac{2}{\sigma^2} (r_x - r_y \cos(\alpha_x - \alpha_y)) K(x, y) \text{ and}$$

$$\begin{aligned} \frac{\partial^2 K}{\partial r_y \partial r_x}(x, y) &= \frac{2}{\sigma^2} \cos(\alpha_x - \alpha_y) K(x, y) \\ &\quad + \frac{4}{\sigma^4} (r_x - r_y \cos(\alpha_x - \alpha_y)) (r_y - r_x \cos(\alpha_x - \alpha_y)) K(x, y). \end{aligned}$$

7.2.2. *Spectral decomposition.* — Using parity arguments, we obtain that the eigenvalues of \mathcal{L}_D are, for $n \geq 0$,

$$\lambda_n^D = R e^{-\frac{2R^2}{\sigma^2}} \int_0^{2\pi} e^{\frac{2R^2}{\sigma^2} \cos \theta} \cos(n\theta) d\theta.$$

The ones of \mathcal{L}_N are, for $n \geq 0$,

$$\lambda_n^N = \int_0^{2\pi} [A \cos \theta + B(1 + \cos^2 \theta)] e^{-\frac{2R^2}{\sigma^2} (1 - \cos \theta)} \cos(n\theta) R d\theta,$$

with $A = \frac{2}{\sigma^2} - \frac{8R^2}{\sigma^4}$ and $B = \frac{4R^2}{\sigma^4}$.

The two operators \mathcal{L}_D and \mathcal{L}_N admit the same eigenfunctions. λ_0^D and λ_0^N are of *multiplicity 1* and associated with

$$(7.3) \quad \begin{array}{ccc} \tilde{\phi}_0 : & [0, 2\pi) & \longrightarrow \mathbb{R} \\ & \alpha & \longmapsto \frac{1}{\sqrt{2\pi R}} \end{array}$$

For $n \geq 1$, λ_n^D and λ_n^N are of *multiplicity 2* and associated with, for $\alpha \in [0, 2\pi)$,

$$(7.4) \quad \tilde{\phi}_n^c(\alpha) = \frac{1}{\sqrt{\pi R}} \cos n\alpha \text{ and } \tilde{\phi}_n^s(\alpha) = \frac{1}{\sqrt{\pi R}} \sin n\alpha.$$

Remark 7.2. — The two spaces $\overline{\mathfrak{F}_D(\mathcal{H})}^{L^2}$ and $\overline{\mathfrak{F}_N(\mathcal{H})}^{L^2}$ are the same and correspond to the linear subspace of 2π -periodic functions of $L_{loc}^2(\mathbb{R})$ (restricted to $[0, 2\pi)$). As the set of all eigenfunctions of each operator \mathcal{L}_D and \mathcal{L}_N coincides with the classical discrete Fourier basis, \mathcal{L}_D and \mathcal{L}_N do not admit other non-null eigenvalue.

Concerning the operator \mathcal{L}_D , one can for instance consult [MNY06] where similar spectral problems are studied (see also Remark 7.5).

We are now able to express the orthonormal bases of \mathcal{H}_{M_D} and \mathcal{H}_{M_N} , associated with \mathfrak{F}_D and \mathfrak{F}_N respectively (see Proposition 3.8). For \mathfrak{F}_D , we introduce the elements ϕ_0^D , ϕ_n^{cD} and $\phi_n^{sD} \in \mathcal{H}_{M_D}$, $n \geq 1$, with for instance

$$(7.5) \quad \forall n \geq 1, \forall x \in \mathcal{X}, \phi_n^{sD}(x) = \frac{1}{\lambda_n^D} \int_0^{2\pi} K(s_{R,\theta}, x) \frac{\sin(n\theta)}{\sqrt{\pi R}} R d\theta.$$

For \mathfrak{F}_N , we introduce ϕ_0^N , ϕ_n^{cN} and $\phi_n^{sN} \in \mathcal{H}_{M_N}$, $n \geq 1$, with

$$(7.6) \quad \forall n \geq 1, \forall x \in \mathcal{X}, \phi_n^{cN}(x) = \frac{1}{\lambda_n^N} \int_0^{2\pi} \frac{\partial K}{\partial r_s}(s_{R,\theta}, x) \frac{\cos(n\theta)}{\sqrt{\pi R}} R d\theta.$$

Examples of numerical computations are presented in Section 7.5. We are now going to study the behavior on the circle \mathcal{C} of this two families of functions.

7.3. An interesting operator. — For $x_{R,\alpha} \in \mathcal{C}$, let us consider the values

$$\langle \phi_0^N, \delta_{x_{R,\alpha}} \rangle_{E,E'}, \langle \phi_n^{cN}, \delta_{x_{R,\alpha}} \rangle_{E,E'} \text{ and } \langle \phi_n^{sN}, \delta_{x_{R,\alpha}} \rangle_{E,E'} \text{ for } n \geq 1,$$

$$\langle \phi_0^D, \eta_{x_{R,\alpha}} \rangle_{E,E'}, \langle \phi_n^{cD}, \eta_{x_{R,\alpha}} \rangle_{E,E'} \text{ and } \langle \phi_n^{sD}, \eta_{x_{R,\alpha}} \rangle_{E,E'} \text{ for } n \geq 1.$$

It appears that those ones are all linked with an operator J_ν given by,

$$(7.7) \quad \forall f \in L^2([0, 2\pi]), J_\nu[f](\alpha) = \int_0^{2\pi} \frac{\partial K}{\partial r_s}(x_{R,\alpha}, s_{R,\theta}) f(\theta) R d\theta.$$

Indeed, we have for instance, for $n \geq 1$,

$$\langle \phi_n^{cN}, \delta_{x_{R,\alpha}} \rangle_{E,E'} = \frac{1}{\lambda_n^N} \int_0^{2\pi} \frac{\partial K}{\partial r_s}(s_{R,\theta}, x_{R,\alpha}) \frac{\cos(n\theta)}{\sqrt{\pi R}} R d\theta,$$

$$\langle \phi_n^{cD}, \eta_{x_{R,\alpha}} \rangle_{E,E'} = \frac{1}{\lambda_n^D} \int_0^{2\pi} \frac{\partial K}{\partial r_x}(s_{R,\theta}, x_{R,\alpha}) \frac{\cos(n\theta)}{\sqrt{\pi R}} R d\theta$$

and $\frac{\partial K}{\partial r_s}(s_{R,\theta}, x_{R,\alpha}) = \frac{\partial K}{\partial r_x}(s_{R,\theta}, x_{R,\alpha})$. The operator J_ν is *self-adjoint but not positive*. For, $n \geq 0$, its eigenvalues are

$$\rho_n = \int_0^{2\pi} \frac{-2R}{\sigma^2} (1 - \cos \theta) e^{-\frac{2R^2}{\sigma^2}(1 - \cos \theta)} \cos(n\theta) R d\theta.$$

We remark in particular that $\rho_0 < 0$. The eigenvalue ρ_0 is of multiplicity 1 and is associated with the same eigenfunction $\tilde{\phi}_0$ than λ_0^D and λ_0^N . For $n \geq 1$, the ρ_n are of multiplicity 2 and are also associated with the same eigenfunctions $\tilde{\phi}_n^c$ and $\tilde{\phi}_n^s$ than λ_n^D and λ_n^N . The same argument than the one used in Remark 7.2 assures that J_ν does not admit other non-null eigenvalue.

The spectrum of the operator J_ν has a particular behavior since the number of its negative eigenvalues depends of the ratio between R and σ^2 . The values of ρ_n , $0 \leq n \leq 30$, for $R = 3$ and $\sigma^2 = 2$ are presented in Figure 2.

Concerning the orthonormal basis of \mathcal{H}_{M_N} associated with \mathfrak{F}_N , we finally obtain, for $x_{R,\alpha} = (R \cos \alpha, R \sin \alpha) \in \mathcal{C}$,

$$(7.8) \quad \begin{aligned} \langle \phi_0^N, \delta_{x_{R,\alpha}} \rangle_{E,E'} &= \frac{\rho_0}{\lambda_0^N} \tilde{\phi}_0(\alpha) \text{ and, for all } n \geq 1, \\ \langle \phi_n^{cN}, \delta_{x_{R,\alpha}} \rangle_{E,E'} &= \frac{\rho_n}{\lambda_n^N} \tilde{\phi}_n^c(\alpha) \text{ and } \langle \phi_n^{sN}, \delta_{x_{R,\alpha}} \rangle_{E,E'} = \frac{\rho_n}{\lambda_n^N} \tilde{\phi}_n^s(\alpha). \end{aligned}$$

For the one of \mathcal{H}_{M_N} associated with \mathfrak{F}_N , we find

$$(7.9) \quad \begin{aligned} \langle \phi_0^D, \eta_{x_{R,\alpha}} \rangle_{E,E'} &= \frac{\rho_0}{\lambda_0^D} \tilde{\phi}_0(\alpha) \text{ and, for all } n \geq 1, \\ \langle \phi_n^{cD}, \eta_{x_{R,\alpha}} \rangle_{E,E'} &= \frac{\rho_n}{\lambda_n^D} \tilde{\phi}_n^c(\alpha) \text{ and } \langle \phi_n^{sD}, \eta_{x_{R,\alpha}} \rangle_{E,E'} = \frac{\rho_n}{\lambda_n^D} \tilde{\phi}_n^s(\alpha). \end{aligned}$$

7.4. Double Constraint. — We combine the results of the two preceding sections in order to obtain a model that both takes account of the values of the function and of its radial derivative on \mathcal{C} . We present an original way to express the kernel $K_{0_C}(\cdot, \cdot)$ of the subspace \mathcal{H}_{0_C} of functions $h \in \mathcal{H}$ such that,

$$\forall t \in \mathcal{C}, \langle h, \eta_t \rangle_{E,E'} = 0 \text{ and } \langle h, \delta_t \rangle_{E,E'} = 0.$$

We use Section 7.2 in order to describe the kernels $T_{\mathcal{H}_{0_N}}$ of \mathcal{H}_{0_N} and $T_{\mathcal{H}_{0_D}}$ of \mathcal{H}_{0_D} . We next consider the interpolation problem associated with $T_{\mathcal{H}_{0_N}}$ and M_D (arbitrary choice, we could equivalently consider the problem associated with $T_{\mathcal{H}_{0_D}}$ and M_N , see Remark 7.4).

Remark 7.3. — This two approaches consist in considering the two decompositions $P_{\mathcal{H}_{0_C}} = P_{\mathcal{H}_{0_D}} P_{\mathcal{H}_{0_N}}$ or $P_{\mathcal{H}_{0_C}} = P_{\mathcal{H}_{0_N}} P_{\mathcal{H}_{0_D}}$, where $P_{\mathcal{H}_{0_C}}$, $P_{\mathcal{H}_{0_D}}$ and $P_{\mathcal{H}_{0_N}}$ are the orthogonal projections of \mathcal{H} onto \mathcal{H}_{0_C} , \mathcal{H}_{0_D} and \mathcal{H}_{0_N} respectively.

We consider the kernel $K_{0_N}(\cdot, \cdot)$ of \mathcal{H}_{0_N} , the subspace of functions $h \in \mathcal{H}$ such that $\frac{\partial h}{\partial r_t}(t) = 0$, for all $t \in \mathcal{C}$. We recall that

$$\begin{aligned} K_{0_N}(x, y) &= K(x, y) - K_{M_N}(x, y) \text{ with} \\ K_{M_N}(x, y) &= \lambda_0^N \phi_0^N(x) \phi_0^N(y) + \sum_{n \geq 1} \lambda_n^N [\phi_n^{cN}(x) \phi_n^{cN}(y) + \phi_n^{sN}(x) \phi_n^{sN}(y)]. \end{aligned}$$

Thanks to the parameterization γ_D and the kernel $K_{0_N}(\cdot, \cdot)$, we define a regular embedding of $\mathcal{H}_{M_N} \cap \mathcal{H}_{0_N}$ into $L^2([0, 2\pi])$. We finally obtain the integral operator \mathcal{L}_{C1} defined by, for $f \in L^2([0, 2\pi])$ and $\alpha \in [0, 2\pi)$,

$$\mathcal{L}_{C1}[f](\alpha) = \int_0^{2\pi} K_{0_N}(x_{R,\alpha}, s_{R,\theta}) f(\theta) R d\theta.$$

From the study of the operator J_ν (see Section 7.3), we obtain that the eigenvalues λ_n^{C1} , $n \in \mathbb{N}$, of \mathcal{L}_{C1} are given by

$$(7.10) \quad \lambda_n^{C1} = \lambda_n^D - \frac{\rho_n^2}{\lambda_n^N}.$$

The eigenvalue λ_0^{C1} is associated with the eigenfunction $\tilde{\phi}_0$ of equation (7.3). For $n \geq 1$, λ_n^{C1} is of multiplicity 2 and associated with $\tilde{\phi}_n^c$ and $\tilde{\phi}_n^s$ (equation (7.4)).

We finally introduce the elements ϕ_0^{C1} , ϕ_n^{cC1} and ϕ_n^{sC1} , $n \geq 1$, which are associated with \mathcal{L}_ν^{C1} via Proposition 3.8. Straightforward calculations give

$$(7.11) \quad \begin{aligned} \phi_0^{C1} &= \frac{1}{\lambda_0^{C1}} (\lambda_0^D \phi_0^D - \rho_0 \phi_0^N) \text{ and, for } n \geq 1, \\ \phi_n^{cC1} &= \frac{1}{\lambda_n^{C1}} (\lambda_n^D \phi_n^{cD} - \rho_n \phi_n^{cN}) \text{ and } \phi_n^{sC1} = \frac{1}{\lambda_n^{C1}} (\lambda_n^D \phi_n^{sD} - \rho_n \phi_n^{sN}). \end{aligned}$$

Remark 7.4. — Instead of first considering M_N and next M_D , one can operate in an inverse way. This leads to the study of the operator

$$\mathcal{L}_{C2}[f](\alpha) = \int_0^{2\pi} \frac{\partial^2 K_{0D}}{\partial r_s \partial r_x} (x_{R,\alpha}, s_{R,\theta}) f(\theta) R d\theta,$$

which is associated with $K_{0D}(\cdot, \cdot)$ and M_N via the parameterization γ_N . Using the results of Section 7.3, we find that the eigenvalues λ_n^{C2} , $n \in \mathbb{N}$, of \mathcal{L}_{C2} are

$$(7.12) \quad \lambda_n^{C2} = \lambda_n^N - \frac{\rho_n^2}{\lambda_n^D}.$$

λ_0^{C2} is of multiplicity 1 and is associated with $\tilde{\phi}_0$. In a same way λ_n^{C2} , for $n \geq 1$, is of multiplicity 2 and is associated with $\tilde{\phi}_n^c$ and $\tilde{\phi}_n^s$. We finally obtain

$$(7.13) \quad \begin{aligned} \phi_0^{C2} &= \frac{1}{\lambda_0^{C2}} (\lambda_0^N \phi_0^N - \rho_0 \phi_0^D) \text{ and for } n \geq 1, \\ \phi_n^{cC2} &= \frac{1}{\lambda_n^{C2}} (\lambda_n^N \phi_n^{cN} - \rho_n \phi_n^{cD}) \text{ and } \phi_n^{sC2} = \frac{1}{\lambda_n^{C2}} (\lambda_n^N \phi_n^{sN} - \rho_n \phi_n^{sD}). \end{aligned}$$

Remark 7.5. — If one conserves the same parameterizations γ_D and γ_N and the same auxiliary space $L^2([0, 2\pi])$, one can in fact obtain similar results for any stationary covariance kernel with required regularity (in the sense that it defines a Hilbert subspace of $E = \mathcal{E}^1$) and of the type

$$K(x, y) = \int_{\mathbb{R}} e^{-i\xi \|x-y\|} d\kappa(\xi),$$

with $\|\cdot\|$ the euclidean norm of \mathbb{R}^2 (and $i^2 = -1$) and where κ is a finite symmetric positive measure on \mathbb{R} (see the Bochner's Theorem, for instance in [CS02]). In particular, from the same arguments of parity than the ones leading to the spectral decomposition of the operators \mathcal{L}_D , \mathcal{L}_N and J_ν of Sections 7.2 and 7.3, we deduce that the eigenfunctions of the integral operators associated with such a kernel $K(\cdot, \cdot)$ are still the same functions $\tilde{\phi}_0$, $\tilde{\phi}_n^c$ and $\tilde{\phi}_n^s$, for $n \geq 1$, of equations (7.3) and (7.4) (discrete Fourier basis). Only the eigenvalues change.

7.5. Numerical application. — In this last section, we compute some of the involved quantities for $R = 3$ and $\sigma^2 = 2$. All computations have been performed with the free software R [R D08]. In particular, the implied integrals have been evaluated by quadrature (rectangle method).

7.5.1. Spectral computations. — We first compute the eigenvalues λ_n^D , λ_n^N and ρ_n of the operators \mathcal{L}_D , \mathcal{L}_N and J_ν , for $0 \leq n \leq 30$. Using equations (7.10) and (7.12), we then directly obtain the values of λ_n^{C1} and λ_n^{C2} . The results are listed in Figure 2 (we do not represent the eigenvalues λ_n^{C2}).




FIGURE 2. Eigenvalues λ_n^D , λ_n^N , ρ_n and λ_n^{C1} , for $0 \leq n \leq 30$, $R = 3$ and $\sigma^2 = 2$.

The second step consists in the computation, thanks to expressions (7.5) and (7.6), of the orthonormal bases of \mathcal{H}_{M_D} and \mathcal{H}_{M_N} associated with \mathfrak{F}_D and \mathfrak{F}_N respectively.

From equation (7.11) and (7.13), we then directly obtain the elements associated with the operators \mathcal{L}_{C1} and \mathcal{L}_{C2} . Examples are given in Figure 3. As expected, ϕ_1^{sC2} vanishes on the circle \mathcal{C} and the radial derivative of ϕ_8^{cC1} also vanishes on \mathcal{C} .

7.5.2. Approximations by truncation. — We now consider approximation by truncation of the different involved kernels. In each case, we conserve the terms which are associated with the most important eigenvalues. We use the same notations and numberings that in Sections 7.2 and 7.4. For $l \in \mathbb{N}^*$, we introduce the following truncated kernels (associated with M_D and M_N),

$$K_{0_D}^{trc}(x, y) = K(x, y) - K_{M_D}^{trc}(x, y), \text{ where}$$




FIGURE 3. Graphs of ϕ_0^D (top-left), ϕ_4^{cN} (top-right), ϕ_1^{sC2} (bottom-left) and ϕ_8^{cC1} (bottom-right) on $[-5, 5]^2$, for $R = 3$ and $\sigma^2 = 2$.

$$K_{M_D}^{trc}(x, y) = \lambda_0^D \phi_0^D(x) \phi_0^D(y) + \sum_{n=1}^l \lambda_n^D [\phi_n^{cD}(x) \phi_n^{cD}(y) + \phi_n^{sD}(x) \phi_n^{sD}(y)] \text{ and}$$

$$K_{0_N}^{trc}(x, y) = K(x, y) - K_{M_N}^{trc}(x, y), \text{ where}$$

$$K_{M_N}^{trc}(x, y) = \lambda_0^N \phi_0^N(x) \phi_0^N(y) + \sum_{n=1}^l \lambda_n^N [\phi_n^{cN}(x) \phi_n^{cN}(y) + \phi_n^{sN}(x) \phi_n^{sN}(y)].$$

Such a kernel $K_{M_N}^{trc}(\cdot, \cdot)$ is hence the approximation by truncation of $K_{M_N}(\cdot, \cdot)$ based on the $N_{trc} = 2l + 1$ largest eigenvalues of the operator \mathcal{L}_N . Remark that for each

eigenvalue (of multiplicity 2) λ_n^N , $1 \leq n \leq l$, we have chosen to consider the both associated eigenvectors ϕ_n^{cN} and ϕ_n^{sN} .

We finally define the approximated kernel $K_{0_{C1}}^{trc}(\cdot, \cdot)$ given by,

$$\begin{aligned} \forall x \text{ and } y \in \mathcal{X}, K_{0_{C1}}^{trc}(x, y) &= K_{0_N}^{trc}(x, y) - \lambda_0^{C1} \phi_0^{C1}(x) \phi_0^{C1}(y) \\ &\quad - \sum_{n=1}^l \lambda_n^{C1} [\phi_n^{cC1}(x) \phi_n^{cC1}(y) + \phi_n^{sC1}(x) \phi_n^{sC1}(y)]. \end{aligned}$$

We fix $l = 15$ and compute the kernel $K_{0_{C1}}^{trc}(\cdot, \cdot)$. This approximated kernel is hence based on the $N_{trc} = 31$ largest eigenvalues of the operators \mathcal{L}_N and \mathcal{L}_{C1} . Figure 4 shows the sample path of a centered Gaussian process with covariance $K_{0_{C1}}^{trc}(\cdot, \cdot)$ and the graph of the function $x \mapsto K_{0_{C1}}^{trc}(x, x)$.




FIGURE 4. Sample path of a centered Gaussian process with covariance $K_{0_{C1}}^{trc}(\cdot, \cdot)$ (left) and Graphical representation of $x \mapsto K_{0_{C1}}^{trc}(x, x)$ on $[-4, 4]^2$ (right), for $R = 3$, $\sigma^2 = 2$ and $N_{trc} = 31$.

The kernel $K_{0_C}(\cdot, \cdot) = K_{0_{C1}}(\cdot, \cdot)$ vanishes on \mathcal{C} (in particular $K_{0_{C1}}(t, t) = 0$ for all $t \in \mathcal{C}$). Concerning its approximation $K_{0_{C1}}^{trc}(\cdot, \cdot)$, we deduce from Sections 7.3 and 7.4 that, for all $t \in \mathcal{C}$, the value of $K_{0_{C1}}^{trc}(t, t)$ is *constant* and

$$\begin{aligned} K_{0_{C1}}^{trc}(t, t) = K_{0_D}^{trc}(t, t) &= K_{0_N}^{trc}(t, t) - \frac{1}{2\pi R} \left(\lambda_0^{C1} + 2 \sum_{n=0}^l \lambda_n^{C1} \right) \\ &= 1 - \frac{1}{2\pi R} \left(\lambda_0^D + 2 \sum_{n=1}^l \lambda_n^D \right). \end{aligned}$$

For $l = 15$, we obtain $K_{0_{C1}}^{trc}(t, t) \approx 1.402309\text{e-}06$, for all $t \in \mathcal{C}$. Additional considerations concerning this approximation are given in Section 7.5.3.

For the radial derivative, we also find a constant value, *i.e.* for all $t \in \mathcal{C}$,

$$\begin{aligned} \left. \frac{\partial^2 K_{0_{C1}}^{trc}}{\partial r_x \partial r_y}(x, y) \right|_{x=y=t} &= \left. \frac{\partial^2 K_{0_N}^{trc}}{\partial r_x \partial r_y}(x, y) \right|_{x=y=t} \\ &= \frac{2}{\sigma^2} - \frac{1}{2\pi R} \left(\lambda_0^N + 2 \sum_{n=1}^l \lambda_n^N \right). \end{aligned}$$

For $l = 15$, we obtain $\left. \frac{\partial^2 K_{0_{C1}}^{trc}}{\partial r_x \partial r_y}(x, y) \right|_{x=y=t} \approx 1.483913\text{e-}05$.

7.5.3. Comparison with discretization. — In this last section, we focus on the interpolation problem associated with \mathcal{H} and M_D and compare the approximations obtain by truncation and discretization.

Hence, we consider a set of $N_{dis} \in \mathbb{N}^*$ points $x_{R, \theta_k} = (R \cos \theta_k, R \sin \theta_k) \in \mathcal{C}$ for $k \in \{0, \dots, N-1\}$ uniformly distributed on \mathcal{C} , that is $\theta_k = \frac{2\pi k}{N_{dis}}$. We then introduce the linear subspace of E'

$$M_D^{dis} = \text{span} \left\{ \delta_{x_{R, \theta_k}}, k \in \{0, \dots, N_{dis} - 1\} \right\}.$$

We approximate the interpolation problem associated with M_D by replacing it by M_D^{dis} (let us signalize that it is, up to a rotation, the optimal way to discretize this problem in regards of $\|\cdot\|_{\gamma, \nu}$). This finally leads to the orthogonal decomposition $\mathcal{H} = \mathcal{H}_{M_D}^{dis} + \mathcal{H}_{0_D}^{dis}$ and solutions of this new discretized problem can be computed by the classical approach of Section 5.1. We denote by $K_{0_D}^{dis}(\cdot, \cdot)$ the reproducing kernel of the subspace $\mathcal{H}_{0_D}^{dis}$.

In order to compare the overall quality of the approximations obtained by truncation and discretization, we consider the terms

$$\text{Err}_D^{dis} = \int_0^{2\pi} K_{0_D}^{dis}(s_{R, \theta}, s_{R, \theta}) R d\theta \text{ and } \text{Err}_D^{trc} = \int_0^{2\pi} K_{0_D}^{trc}(s_{R, \theta}, s_{R, \theta}) R d\theta.$$

We remind that this kind of terms naturally arises when one aims at studying, for instance in the discretization case, quantities of the type (see Theorem 4.2 and Proposition 6.4)

$$\left\| \mathbb{P}_{\mathcal{H}_M}[\varphi] - \mathbb{P}_{\mathcal{H}_M^{dis}}[\varphi] \right\|_{\gamma, \nu}^2, \text{ with } \varphi \in \mathcal{H}.$$

Because of the optimality of $K_{0_D}^{trc}(\cdot, \cdot)$ in regards of $\|\cdot\|_{\gamma, \nu}$ (see Section 6.4), for $N_{dis} = N_{trc}$, we always have $\text{Err}_D^{trc} \leq \text{Err}_D^{dis}$ (note that these two terms tend to zero as N_{dis} and N_{trc} tend to $+\infty$, see also Figure 5). For instance, for $R = 3$, $\sigma^2 = 2$ and $N_{dis} = N_{trc} = 31$, we obtain

$$\text{Err}_D^{dis} \approx 4.438046\text{e-}05 \text{ and } \text{Err}_D^{trc} \approx 2.64329\text{e-}05.$$

Although the gain in considering truncation instead of discretization could in this case appear numerically negligible, one has to remark that the behavior on \mathcal{C} of this two kinds of approximation is totally different (see Figure 5). It can hence be said




FIGURE 5. Graphs of $\theta \mapsto K_{0D}^{trc}(s_{R,\theta}, s_{R,\theta})$ and $\theta \mapsto K_{0D}^{dis}(s_{R,\theta}, s_{R,\theta})$ (with $\theta \in [0, 2\pi)$) for $N_{trc} = N_{dis} = 31$ (left) and values of the terms Err_D^{dis} and Err_D^{trc} for $10 \leq N_{dis}, N_{trc} \leq 24$ (right) with $R = 3$ and $\sigma^2 = 2$.

that truncation leads to a global approximation on \mathcal{C} (with respect to $\|\cdot\|_{\gamma,\nu}$) whereas discretization leads to a local approximation (localized at the discretization points).

Acknowledgments

The authors would like to thank the two anonymous reviewers for their valuable comments and suggestions that significantly improved the manuscript and the three associate editors Jean-Marc AZAÏS, Fabrice GAMBOA and Bertrand IOOSS for the work they have put into the organization of this special issue. Many thanks as well to Laurent CARRARO and Mario AHUES for their wise advices and also to Georges GREKOS and Éric CANON for their suggestions on drafting.

References

- [Aro50] N. Aronszajn. Theory of reproducing kernels. *Trans. Amer. Math. Soc.*, 63:337–404, 1950.
- [Att92] M. Atteia. *Hilbertian kernels and spline functions*. North-holland, 1992.
- [Bax76] P. Baxendale. Gaussian measures on function spaces. *American Journal of Mathematics*, 98(4):891–952, 1976.
- [Bou59] N. Bourbaki. *Éléments de mathématique: Chapitre 6, Intégration vectorielle*. Hermann, 1959.
- [BTA04] A. Berlinet and C. Thomas-Agnan. *Reproducing kernel Hilbert spaces in probability and statistics*. Springer Netherlands, 2004.

- [CS02] F. Cucker and S. Smale. On the mathematical foundations of learning. *Bulletin (new series) of the American Mathematical Society*, 39(1):1–49, 2002.
- [DFLC71] R.M. Dudley, J. Feldman, and L. Le Cam. On Seminorms and Probabilities, and Abstract Wiener Spaces. *Annals of Mathematics*, 93(2):390–408, 1971.
- [Dud10] R.M. Dudley. *Sample functions of the Gaussian process*. Springer, 2010.
- [For85] R.M. Fortet. Les operateurs intégraux dont le noyau est une covariance. *Trabajos de Estadística y de Investigación Operativa*, 36:133–144, 1985.
- [Gau11] B. Gauthier. *Approche spectrale pour l’interpolation à noyaux et positivité conditionnelle*. PhD thesis, Ecole des Mines de Saint-Etienne, 2011.
- [Gro67] L. Gross. Abstract Wiener spaces. *Proc. Fifth Berkeley Symp. on Math. Statist. and Prob.*, 2:31–42, 1967.
- [Jan97] S. Janson. *Gaussian Hilbert Spaces*. Cambridge University Press, 1997.
- [Kue71] J. Kuelbs. Expansions of vectors in a Banach space related to Gaussian measures. *Proceedings of the American Mathematical Society*, 27(2):364–370, 1971.
- [MNY06] H.Q. Minh, P. Niyogi, and Y. Yao. *Learning Theory*, chapter Mercer’s Theorem, Feature Maps and Smoothing, pages 154–168. Springer Berlin / Heidelberg, 2006.
- [NW74] M.Z. Nashed and G. Wahba. Generalized inverses in reproducing kernel spaces: An approach to regularization of linear operator equations. *SIAM J. Math. Anal.*, 5(6):974–987, 1974.
- [Par62] E. Parzen. Extraction and detection problems and reproducing kernel hilbert spaces. *J. Soc. Ind. Appl. Math., Ser. A, Control*, 1:35–62, 1962.
- [R D08] R Development Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2008. ISBN 3-900051-07-0.
- [Raj72] B.S. Rajput. On Gaussian measures in certain locally convex spaces. *Journal of Multivariate Analysis*, 2(3):282–306, 1972.
- [Rud95] W. Rudin. *Analyse fonctionnelle*. Ediscience International, 1995.
- [RW06] C.E Rasmussen and C.K.I. Williams. *Gaussian Processes for Machine Learning*. The MIT Press, 2006.
- [Sch64] L. Schwartz. Sous-espaces hilbertiens d’espaces vectoriels topologiques et noyaux associés. *J. Anal. Math.*, 13:115–256, 1964.
- [Sch79] L. Schwartz. *Analyse Hilbertienne*. Hermann, 1979.
- [Ste99] M.L. Stein. *Interpolation of Spatial Data: some theory for kriging*. Springer Verlag, 1999.
- [Tal83] M. Talagrand. Mesures gaussiennes sur un espace localement convexe. *Probability Theory and Related Fields*, 64(2):181–209, 1983.
- [TV07] V. Tarieladze and N. Vakhania. Disintegration of Gaussian measures and average-case optimal algorithms. *Journal of Complexity*, 23(4-6):851–866, 2007.
- [Wah90] G. Wahba. *Spline Models for Observational Data*. SIAM, 1990.
- [Wal67] J.B. Walsh. A note on uniform convergence of stochastic processes. In *Proc. Amer. Math. Soc.*, volume 18, pages 129–132, 1967.

B. GAUTHIER, Université Jean-Monnet de Saint-Étienne, ICJ, URM 5208, PRES univ. de Lyon
E-mail : bertrand.gauthier.314@gmail.com

X. BAY, École des Mines de Saint-Étienne, Institut Fayol
E-mail : bay@emse.fr